

Triple Nitrate Isotopes Indicate Differing Nitrate Source Contributions to Streams Across a Nitrogen Saturation Gradient

Lucy A. Rose,^{1*} Emily M. Elliott,¹ and Mary Beth Adams²

¹Department of Geology and Planetary Science, University of Pittsburgh, 4107 O'Hara Street, 200 Space Research and Coordination Center, Pittsburgh, Pennsylvania 15260, USA; ²Fernow Experimental Forest, USDA Forest Service, 180 Canfield Street, Morgantown, West Virginia 26505, USA

ABSTRACT

Nitrogen (N) deposition affects forest biogeochemical cycles worldwide, often contributing to N saturation. Using long-term (>30-year) records of stream nitrate (NO_3^-) concentrations at Fernow Experimental Forest (West Virginia, USA), we classified four watersheds into N saturation stages ranging from Stage 0 (N-limited) to Stage 3 (N-saturated). We quantified NO_3^- contributions from atmospheric and microbial sources using $\delta^{15}\text{N}$, $\delta^{18}\text{O}$, and $\Delta^{17}\text{O}$ of NO_3^- and characterized the concentrations and isotopes of NO_3^- in precipitation. Despite receiving identical atmospheric inputs, the proportions of atmospheric NO_3^- in streams averaged from 7 to 10% in the hardwood watersheds (stages 1, 2, and 3) and 54% in the conifer watershed (Stage 0). This suggests that the hardwood watersheds may be less responsive to future reductions in N deposition than the conifer watershed, at least in the short term. As shown in other studies, atmospheric NO_3^- proportions were

higher during stormflow. Despite large proportions of atmospheric NO_3^- in the Stage 0 stream, total atmospheric NO_3^- -N flux from this watershed (2.9 g ha^{-1}) was lower than fluxes in the other watersheds (range = 117.8 – 338.5 g ha^{-1}). Seasonal patterns of $\delta^{15}\text{N}$ - NO_3^- in the hardwood watersheds suggest enrichment of the soil NO_3^- pool during the growing season due to plant uptake. In all watersheds, $\delta^{18}\text{O}$ -based mixing models over-estimated atmospheric NO_3^- contributions to streams by up to 12% compared to $\Delta^{17}\text{O}$ -based estimates. Our results highlight the importance of atmospheric deposition as a NO_3^- source in low-concentration streams and demonstrate the advantage of using $\Delta^{17}\text{O}$ - NO_3^- over $\delta^{18}\text{O}$ - NO_3^- for NO_3^- source apportionment.

Key words: nitrogen saturation; nitrogen deposition; nitrate isotopes; Fernow Experimental Forest; $\delta^{15}\text{N}$ - NO_3^- ; $\delta^{18}\text{O}$ - NO_3^- ; $\Delta^{17}\text{O}$ - NO_3^- .

Received 12 July 2014; accepted 1 May 2015

Author contributions Emily M. Elliott designed the study. Lucy A. Rose executed the field work and data analysis. Lucy A. Rose, Emily M. Elliott, and Mary Beth Adams collaborated on the preparation of the paper.

*Corresponding author; e-mail: lab125@pitt.edu

INTRODUCTION

Long-term elevated deposition of atmospheric nitrate (NO_3^-) and ammonium (NH_4^+) to forests can lead to nitrogen saturation (Peterjohn and others 1996; Gilliam and others 2001; Aber and others 2003), with ecosystem responses including increased nitrification rates (Peterjohn and others

1996; Gilliam and others 2001; Lovett and Goodale 2011), decreased soil fertility (Adams and others 2007), and increased NO_3^- and base cation leaching from soils leading to soil acidification (Adams and others 2006; Fernandez and others 2010). Nitrate concentrations and fluxes in soil and stream waters have been previously used to assess the N saturation status of forested watersheds (Stoddard 1994; Peterjohn and others 1996; Aber and others 2003; Lovett and Goodale 2011). However, ecosystem responses to atmospheric deposition can vary depending on site factors such as land-use and disturbance histories, species composition, and hydrologic regime (Aber and others 2003; Adams and others 2014). When examined at the watershed scale, stream NO_3^- dynamics integrate myriad ecosystem processes; it is therefore important to understand the relative contributions of atmospheric and microbial sources to stream NO_3^- export. As anthropogenic additions to the global N cycle will likely continue unabated for the foreseeable future (Galloway and others 2003), clarifying how post-depositional processes in forests influence the transport and fate of atmospheric NO_3^- is critical to understanding ecosystem responses to chronic N deposition and the development of N saturation.

Aber and others (1989, 1998) described a conceptual model of ecosystem N saturation consisting of four stages, ranging from N-limited (Stage 0) to severely N-saturated (Stage 3); increasing nitrification rates and long-term increases in stream NO_3^- concentrations are important indicators of N saturation status in this conceptual model. Although rigorous testing of the N saturation concept requires extensive characterization of ecosystem pools and fluxes (Lovett and Goodale 2011), long-term (that is, decadal) stream chemistry dynamics can provide a convenient framework for investigating ecosystem responses to atmospheric NO_3^- deposition (Stoddard 1994). When coupled with stable isotope measurements of stream NO_3^- ($\delta^{15}\text{N}$, $\delta^{18}\text{O}$, and $\Delta^{17}\text{O}$), the relative contributions of atmospheric and microbial sources to stream NO_3^- across a range of N saturation stages can be elucidated.

Previous studies have used nitrogen and oxygen stable isotopes to track spatio-temporal changes in atmospheric and microbial NO_3^- pools and to differentiate between them (Durka and others 1994; Burns and Kendall 2002; Pardo and others 2004; Barnes and others 2008; Sebestyen and others 2008, 2014; Goodale and others 2009). For example, temporal patterns in $\delta^{15}\text{N}$ values in precipitation and stream NO_3^- have been used to identify

NO_3^- uptake in forests (Barnes and others 2008). Similarly, $\delta^{18}\text{O}-\text{NO}_3^-$ values have been used to distinguish atmospheric and soil sources and to determine their relative contributions to ecosystem NO_3^- pools (Williard and others 2001; Pardo and others 2004; Goodale and others 2009; Sebestyen and others 2014). However, preferential uptake of lighter isotopes (for example, ^{14}N and ^{16}O) during biological processes such as denitrification can alter $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ values of residual NO_3^- pools, leading to imprecise source apportionment in some cases (Michalski and others 2004; Riha and others 2014). Nitrate isotopic source signatures can also result from processes not dependent on mass; these mass-independent signatures—such as $\Delta^{17}\text{O}-\text{NO}_3^-$ —are not fractionated by biological processes and therefore serve as conservative tracers of atmospheric and microbial NO_3^- (Michalski and others 2004). Because atmospheric NO_3^- is characterized by positive $\Delta^{17}\text{O}$ values and microbial NO_3^- by values of zero or less, $\Delta^{17}\text{O}$ is a robust tool for quantifying the contributions of atmospheric and microbial NO_3^- to surface waters.

Nitrogen cycling is an important, but poorly understood phenomenon in forests, and the importance of atmospheric deposition in determining N exports is still not well elucidated. In this study, we quantified atmospheric and microbial source contributions to stream NO_3^- using triple NO_3^- isotopes ($\delta^{15}\text{N}$, $\delta^{18}\text{O}$, and $\Delta^{17}\text{O}$) of precipitation and stream water in four nearly adjacent watersheds at Fernow Experimental Forest (Figure 1). These watersheds have historically received some of the highest rates of atmospheric NO_3^- deposition in the U.S. (Adams and others 2003). Despite receiving identical atmospheric N inputs, these watersheds demonstrate differing long-term (>30-year) patterns in stream NO_3^- concentrations due to differences in internal N retention (Figure 2). Such stream NO_3^- concentration dynamics are consistent with the progressive stages of watershed N saturation described by Stoddard (1994) and Aber and others (1989, 1998) and are likely influenced to some extent by differences in tree species composition among the watersheds. These decades-long differences in stream NO_3^- concentrations served as the framework for the N saturation gradient examined in this study. Using triple NO_3^- isotope analyses, our study objectives were to (1) characterize and quantify the source contributions to stream NO_3^- in watersheds receiving identical N deposition inputs but exhibiting long-term differences in stream NO_3^- concentrations and (2) elucidate the roles of biological and hydrologic drivers in regulating atmospheric NO_3^- export. Based on the results of prior

studies (for example, Durka and others (1994)), we hypothesized that watersheds with higher N saturation status would have greater proportions of atmospheric NO_3^- in stream water than those with lower N saturation status.

METHODS

Study Site

Fernow Experimental Forest is located in West Virginia, on the unglaciated Allegheny Plateau (Figure 1). Elevations in the study watersheds range from 720 to 865 m, and slopes average about 20%. Bedrock is composed of hard sandstone and softer shale; little water storage occurs in these strata (Reinhart and others 1963; Kochenderfer 2007). Soils are well-drained silt loams of the Calvin series (loamy-skeletal, mixed active, mesic typic Dystrudept), averaging 1 m in depth (Kochenderfer 2007). Infiltration rates are high and most precipitation reaches the streams via subsurface flow (Reinhart and others 1963). During high-intensity rain events, runoff is high and falls off quickly during periods of low-intensity or no pre-

cipitation (Reinhart and others 1963). The growing season extends from late April through October and precipitation is evenly distributed throughout the year, averaging 1450 mm; significant snowpack does not accumulate over long periods. Mixed hardwoods are the dominant forest type at Fernow, with northern red oak (*Quercus rubra*), sugar maple (*Acer saccharum*), red maple (*Acer rubrum*), black cherry (*Prunus serotina*), and yellow poplar (*Liriodendron tulipifera*), the most abundant species. As an exception to these native hardwoods, one research watershed was planted to Norway spruce (*Picea abies*) monoculture in 1973.

The research watersheds for this study were selected based on long-term (>30-year) differences in patterns of mean stream NO_3^- -N concentrations (Figure 2); these long-term differences were used to classify the watersheds into different N saturation stages, ranging from Stage 0 (N-limited) to Stage 3 (severely N-saturated), following the conceptual model of N saturation described by Aber and others (1989). We sampled three mixed hardwood-dominated watersheds (Stages 1, 2, and 3) and one conifer-dominated watershed (Norway




Figure 1. Map of study watersheds within the larger boundary of Fernow Experimental Forest (FEF). Numbers indicate N saturation stage; black dots show weir locations in study watersheds. Contour spacing is 30 m. Inset shows the location of FEF (black dot) in the state of West Virginia (shown in grey).




Figure 2. Monthly stream NO_3^- -N concentrations in the study watersheds at Fernow Experimental Forest averaged from weekly measurements collected from 1983 to 2007. The *four watersheds* exhibit differing long-term concentrations and seasonal patterns of NO_3^- -N in streams; watershed N saturation stage classification was based on these long-term differences in stream NO_3^- -N dynamics.

spruce; Stage 0) for this study. Several site characteristics are similar among the watersheds (for example atmospheric deposition inputs, climate, discharge patterns, geology, soils, elevation) but the watersheds differ with respect to a number of factors including stream NO_3^- -N concentrations, mean streambed and watershed slope, land-use history, and dominant overstorey composition (Table 1). In addition, the riparian zone in the conifer-dominated Stage 0 watershed is characterized by patches of *Sphagnum* sp. in some areas; such features are absent in any of the hardwood-dominated study watersheds. The presence of *Sphagnum* sp. patches in the Stage 0 watershed suggests that riparian wetland areas may be starting to form (Tiner 1993).

Sample Collection

From January through December 2010, 1 L stream samples were collected in acid-washed HDPE bottles from the stream just above a 120° V-notch weir at the base of each watershed. Samples were collected on a weekly basis when streams were flowing. Samples were vacuum-filtered through 0.22 μm polyethersulfone membrane filters to remove suspended solids and biological material. All samples were processed at the US Forest Service Timber and Watershed Laboratory in Parsons, West Virginia within 24 h of collection. Filtered samples were frozen and transported to the University of Pittsburgh, where they remained frozen until isotopic analysis. Weekly samples of wet-only depo-

sition were collected at a National Atmospheric Deposition Program (NADP) National Trends Network site approximately 2 km from the study watersheds from February through December 2010. These samples were filtered through 0.45 μm polyethersulfone membrane filters and archived at 4°C at the NADP Central Analytical Laboratory in Champaign, IL, USA. Precipitation samples were shipped to the University of Pittsburgh and frozen until isotopic analysis.

Isotopic Analysis

Nitrate concentrations were measured by ion chromatography (Dionex ICS-2000) at the University of Pittsburgh. For isotopic analysis, a denitrifying bacteria, *Pseudomonas aureofaciens*, was used to convert aqueous NO_3^- into gaseous N_2O which was introduced into the mass spectrometer (Sigman and others 2001; Casciotti and others 2002). For $\Delta^{17}\text{O}$ analysis, this N_2O was thermally decomposed at 800°C into N_2 and O_2 prior to isotopic analysis following the method described by Kaiser and others (2007). Duplicate samples were analyzed for $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ - NO_3^- (and separately for $\Delta^{17}\text{O}$ - NO_3^-) on an Isoprime Trace Gas and Gilson GX-271 autosampler coupled with an Isoprime Continuous Flow Isotope Ratio Mass Spectrometer (CF-IRMS) at the *Regional Stable Isotope Laboratory for Earth and Environmental Science Research* at the University of Pittsburgh. Isotope values are reported in parts per thousand relative to NO_3^- standards as follows:

Table 1. Description of Study Watersheds

N sat. stage	Mean stream [NO ₃ ⁻ -N] (mg L ⁻¹) 1980–2010	Size (ha)	Dominant overstory species	Forest age (years)	Aspect	Mean stream channel slope (%)	Mean watershed slope (%)	Treatment history
0	0.21	22	<i>Picea abies</i>	40	S	2.4	23.0	Lower half clearcut 1964; upper half clearcut 1967–68; maintained barren until 1969. Planted <i>Picea abies</i> 1973
1	0.39	37	<i>Acer rubrum</i> , <i>Acer saccharum</i>	Uneven	NE	3.5	30.6	Selective cut (~10% of basal area > 27.9 cm dbh removed every 10 years since 1958)
2	0.77	39	<i>Quercus rubra</i> , <i>Acer rubrum</i>	110	SE	10.9	22.3	Control, no treatment
3	1.15	24	<i>Liriodendron tulipifera</i> , <i>Acer rubrum</i> , <i>Quercus rubra</i>	44	E	7.5	24.8	Upper half clearcut 1963–64; lower half clearcut 1966–67; maintained barren until 1969, then natural regeneration

$$\delta^{15}\text{N}, \delta^{18}\text{O}, \text{ and } \delta^{17}\text{O} (\text{‰}) = \left[\left(\frac{R_{\text{sample}}}{R_{\text{standard}}} \right) - 1 \right] \times 1000 \quad (1)$$

where $R = {}^{15}\text{N}/{}^{14}\text{N}$, ${}^{18}\text{O}/{}^{16}\text{O}$, or ${}^{17}\text{O}/{}^{16}\text{O}$. The mass-independent oxygen isotope anomaly ($\Delta^{17}\text{O}-\text{NO}_3^-$) is likewise reported in parts per thousand and calculated using the equation:

$$\Delta^{17}\text{O} (\text{‰}) = \delta^{17}\text{O} - 0.52 \times \delta^{18}\text{O} \quad (2)$$

Samples with low NO_3^- concentrations were pre-concentrated prior to bacterial conversion to N_2O . For pre-concentration, we calculated the sample volume necessary to obtain a final NO_3^- concentration of 20 nmol (for $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ analysis) or 200 nmol (for $\Delta^{17}\text{O}$ analysis) in a 5 ml sample. Appropriate sample volumes were measured into 10% hydrochloric acid-washed Pyrex or Teflon beakers and placed in a drying oven at 60°C until all liquid was evaporated. The interior of each beaker was then rinsed with 10 ml of 18 MΩ water to reconstitute duplicate samples to the appropriate concentration. Samples were prepared for isotopic analysis following the bacterial denitrifier method as previously described. International reference standards were similarly pre-concentrated for correction of pre-concentrated samples.

$\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ values were corrected using international reference standards USGS-32, USGS-34, USGS-35, and IAEA-N3; USGS-34 and USGS-35 were used to correct $\Delta^{17}\text{O}$ values. These standards were also used to correct for linearity and instrument drift. Standard deviations for international reference standards were 0.2, 0.5, and 0.2‰ for $\delta^{15}\text{N}$, $\delta^{18}\text{O}$, and $\Delta^{17}\text{O}$, respectively.

$\delta^{15}\text{N}-\text{NO}_3^-$ values must be corrected for interference from contributions of ${}^{14}\text{N}{}^{14}\text{N}{}^{17}\text{O}$ to m/z 45 (Coplen and others 2004). These contributions were evaluated following the relationship described in Coplen and others (2004), where a 1‰ increase in $\delta^{15}\text{N}$ corresponds to an 18.8‰ increase in $\Delta^{17}\text{O}$. Corrected $\delta^{15}\text{N}$ values in stream samples were zero to 1.3‰ (mean = 0.02‰) lower than uncorrected values, depending on the mass-independent contribution of $\Delta^{17}\text{O}$ in the sample. Because $\Delta^{17}\text{O}-\text{NO}_3^-$ values were low in most samples and because we could not apply the correction to some samples due to a lack of $\Delta^{17}\text{O}$ data (particularly in the Stage 0 watershed), the $\delta^{15}\text{N}$ values presented here do not include the mass-independent $\Delta^{17}\text{O}$ correction. Although interference-corrected $\delta^{15}\text{N}$ data are slightly lower than those presented here, the temporal and spatial patterns in $\delta^{15}\text{N}$ should not be strongly influenced by the omission of this correction.

End-Member Mixing Analysis

We used a two-end-member isotope mixing model approach (using both $\delta^{18}\text{O}$ and $\Delta^{17}\text{O}$ isotope systems) to calculate the fraction of atmospheric NO_3^- exported from each watershed (equation (3)):

$$f_{\text{atm}} = \frac{\chi_{\text{stream}} - \chi_{\text{nitrification}}}{\chi_{\text{atm}} - \chi_{\text{nitrification}}} \quad (3)$$

where χ is $\delta^{18}\text{O}$ or $\Delta^{17}\text{O}$ of NO_3^- . We used the lowest measured stream NO_3^- $\delta^{18}\text{O}$ (or $\Delta^{17}\text{O}$) value in each watershed as the nitrification end-member (Barnes and others 2008). The lowest $\Delta^{17}\text{O}$ – NO_3^- values in the hardwood watersheds were -1.2 , -1.2 , and -1.4‰ in the Stage 1, 2, and 3 watersheds, respectively. In the Stage 0 watershed, a value of zero was used for the nitrification end-member $\Delta^{17}\text{O}$ value, as all stream samples had positive values. For the atmospheric end-member, we used the volume-weighted mean $\delta^{18}\text{O}$ (or $\Delta^{17}\text{O}$) value of all weekly precipitation samples collected during 2010. We used the volume-weighted mean rather than individual precipitation samples collected at the same time as stream samples because hydrologic transit times have not been characterized for most of the study watersheds, and the isotopic composition of atmospheric NO_3^- in a stream sample is unlikely to correspond to that of the precipitation sample collected during the same week.

Nitrate Flux Calculation

Nitrate–N fluxes were calculated in each watershed on a weekly basis using total weekly discharge (based on continuous flow level measurements taken at the weir) and NO_3^- –N concentrations determined from stream samples collected weekly. Total annual NO_3^- –N flux in each watershed was calculated as the sum of all weekly NO_3^- –N fluxes during the study period. Total annual atmospheric NO_3^- –N fluxes were calculated for the entire study period by multiplying the total annual NO_3^- –N flux in each watershed by the discharge-weighted mean proportion of atmospheric NO_3^- in the stream (based on $\Delta^{17}\text{O}$ – NO_3^-). As the proportion of atmospheric NO_3^- in the Stage 0 watershed was based on only those stream samples with NO_3^- –N concentrations sufficient for $\Delta^{17}\text{O}$ – NO_3^- analysis ($n = 4$), the total atmospheric NO_3^- –N flux calculated for this watershed should be interpreted with caution.

Statistical Analysis

We used analysis of variance (ANOVA) to test for significant differences among watersheds. When significant differences were indicated, we applied Tukey's Honestly Significant Difference test to determine which means were significantly different ($\alpha = 0.05$). The experiment-wise error rate was held at $\alpha = 0.05$ for comparisons among watersheds. To test for differences between growing season (May–October) and dormant season (November–April) means within individual watersheds, we used repeated measures ANOVA. All statistical analyses were conducted using volume-weighted concentrations and concentration-weighted isotope values; therefore, all means are reported as weighted values. In order to include samples with NO_3^- concentrations below the instrument detection limit ($0.01 \text{ mg NO}_3^- \text{ L}^{-1}$) in statistical analyses, NO_3^- concentrations for these samples were set at $0.005 \text{ mg NO}_3^- \text{ L}^{-1}$ (one-half of the instrument detection limit). All statistical analyses were conducted using SAS statistical software (SAS Institute, Inc. 2011).

RESULTS

Nitrate–N Concentration

Intermittent streamflow in all study watersheds prevented sample collection during parts of July through October in the Stage 0, 1, and 2 watersheds, and August through October in the Stage 3 watershed. In addition, low stream NO_3^- –N concentrations in the Stage 0 watershed precluded isotopic analysis of several samples. Stream NO_3^- –N concentrations across all watersheds ranged from below the instrument detection limit to 1.91 mg L^{-1} and precipitation NO_3^- –N concentrations ranged from 0.02 to 0.67 mg L^{-1} (Figure 3; Table 2). Mean NO_3^- –N concentrations were significantly different in all watersheds ($P < 0.01$; Table 2); this was expected, given the long-term patterns in stream NO_3^- –N concentrations in these watersheds (Figure 2) and the nature of the study design (that is, stream NO_3^- concentrations were used to define watershed N saturation status). The Stage 1, 2, and 3 (hardwood) watersheds showed significant seasonal differences in stream NO_3^- –N concentrations, with lower mean values during the growing season (May to October) than the dormant season (November to April) ($P < 0.01$; Table 3).




Figure 3. A Nitrate-N concentration, B $\delta^{15}\text{N}-\text{NO}_3^-$, C Daily total discharge, D $\delta^{18}\text{O}-\text{NO}_3^-$, and E $\Delta^{17}\text{O}-\text{NO}_3^-$ of weekly stream and precipitation samples collected at Fernow during 2010. Months shaded in grey represent the growing season. The dashed circle in E indicates samples collected during the snowmelt event on 23 February.

$\delta^{15}\text{N}$ of Nitrate

Across all watersheds, values of $\delta^{15}\text{N}$ in stream NO_3^- ranged from -2.9‰ to $+5.9\text{‰}$, and precipitation values ranged from -6.0‰ to $+5.1\text{‰}$ (Figure 3; Table 2). Mean $\delta^{15}\text{N}-\text{NO}_3^-$ values were not significantly different among the watersheds or precipitation (Table 2). Seasonally, mean stream $\delta^{15}\text{N}-\text{NO}_3^-$ values were significantly higher during the growing season than the dormant season in the Stage 1, 2, and 3 watersheds ($P < 0.01$; Table 3). The seasonal variability in $\delta^{15}\text{N}-\text{NO}_3^-$ may have contributed to the lower mean $\delta^{15}\text{N}-\text{NO}_3^-$ value in the Stage 0 watershed, as streamflow was absent in this watershed during most of the growing season (when $\delta^{15}\text{N}$ values were highest in the Stage 1, 2, and 3 watersheds; Figure 3). In contrast to the seasonal patterns in the hardwood watersheds, the mean $\delta^{15}\text{N}-\text{NO}_3^-$ value in precipitation was lower

during the growing season than the dormant season, but this difference was not statistically significant (Table 3).

$\delta^{18}\text{O}$ of Nitrate

Across all watersheds, $\delta^{18}\text{O}$ values in stream NO_3^- ranged from -9.4 to $+72.8\text{‰}$, whereas precipitation $\delta^{18}\text{O}-\text{NO}_3^-$ values ranged from $+51.1\text{‰}$ to $+81.7\text{‰}$ (Figure 3; Table 2). The highest mean $\delta^{18}\text{O}-\text{NO}_3^-$ value of stream NO_3^- occurred in the Stage 0 watershed ($+40.4\text{‰}$; Table 2). This mean was significantly higher than those in all other watersheds, but significantly lower than the mean precipitation $\delta^{18}\text{O}-\text{NO}_3^-$ value ($+65.3\text{‰}$) ($P < 0.01$ for all comparisons). In all watersheds and in precipitation, mean $\delta^{18}\text{O}-\text{NO}_3^-$ values were significantly higher during the dormant season than the growing season ($P < 0.02$; Table 3).

Table 2. Weighted Means (SE) and Ranges for Stream NO_3^- -N Concentration and Isotope Values

N sat. stage	$[\text{NO}_3^- \text{-N}]$ (mg L^{-1})		$\delta^{15}\text{N-NO}_3^-$ (‰)		$\delta^{18}\text{O-NO}_3^-$ (‰)		$\Delta^{17}\text{O-NO}_3^-$ (‰)	
	Mean	Range	Mean	Range	Mean	Range	Mean	Range
0	0.00 ^d (0.00)	na to 0.02	+1.0 (0.8)	-2.9 to +3.8	+40.4 ^b (8.3)	-5.3 to +72.8	+11.1 ^b (5.8)	+1.4 to +23.9
1	0.25 ^c (0.01)	0.14 to 0.33	+2.2 (0.2)	+1.0 to +5.0	-1.8 ^c (0.7)	-9.4 to +13.4	+0.3 ^c (0.2)	-1.2 to +5.6
2	0.48 ^b (0.02)	0.29 to 0.79	+2.2 (0.1)	+1.5 to +4.3	-1.0 ^c (0.6)	-7.9 to +6.1	+0.5 ^c (0.2)	-1.2 to +3.2
3	0.84 ^a (0.02)	0.56 to 1.91	+2.9 (0.2)	+1.6 to +5.9	-2.7 ^c (0.4)	-7.5 to +4.0	-0.3 ^c (0.1)	-1.4 to +3.0
Precip	0.17 ^c (0.02)	0.02 to 0.67	-0.1 (0.4)	-6.0 to +5.1	+65.3 ^a (1.1)	+51.1 to +81.7	+21.3 ^a (0.7)	+12.3 to +29.1

Samples with concentrations below the instrument detection limit ($n = 5$) are denoted by na. For statistical analysis, NO_3^- -N concentrations for these samples were set to 0.005 mg L^{-1} (one-half the instrument detection limit). Within a column, means followed by different letters are significantly different at $\alpha = 0.05$. For $\delta^{15}\text{N-NO}_3^-$, none of the means were significantly different.

$\Delta^{17}\text{O}$ of Nitrate

$\Delta^{17}\text{O-NO}_3^-$ values in stream water ranged from -1.4‰ to $+23.9\text{‰}$ across all watersheds (Figure 3; Table 2), with a significantly higher mean value in the Stage 0 watershed than in the other watersheds ($P < 0.01$ for all comparisons; Table 2). However, mean stream $\Delta^{17}\text{O-NO}_3^-$ values were not significantly different when ANOVA was conducted using only data for the four dates on which $\Delta^{17}\text{O}$ values were available in the Stage 0 watershed (in other words, $n = 4$ for each watershed). While this analysis allows for a more balanced comparison of $\Delta^{17}\text{O}$ values across watersheds, the small sample sizes weaken the statistical power of ANOVA, making it difficult to discern whether the mean $\Delta^{17}\text{O}$ value in the Stage 0 watershed is truly different from the others. Precipitation $\Delta^{17}\text{O-NO}_3^-$ values ranged from $+12.3$ to $+29.1\text{‰}$, with a mean of $+20.6\text{‰}$. Mean $\Delta^{17}\text{O-NO}_3^-$ values in the Stage 1 and 2 watersheds and in precipitation were significantly higher during the dormant season than the growing season ($P < 0.04$; Table 3). In the Stage 0 and 3 watersheds, mean $\Delta^{17}\text{O-NO}_3^-$ values were also higher during the dormant season than the growing season, but seasonal differences were not statistically significant. In all watersheds, the highest $\Delta^{17}\text{O-NO}_3^-$ value measured during 2010 occurred on 23 February, coincident with a snow-melt event (Figure 3).

Proportions and Fluxes of Atmospheric Nitrate in Streams

We calculated the proportions of atmospheric NO_3^- in streams using both $\delta^{18}\text{O-NO}_3^-$ and $\Delta^{17}\text{O-NO}_3^-$ values to evaluate differences in estimated atmospheric NO_3^- proportions in streams. Substituting $\delta^{18}\text{O-NO}_3^-$ values into equation (3), the mean proportions of atmospheric NO_3^- in the Stage 0, 1, 2, and 3 watersheds were 76, 14, 12, and 9%, respectively. When $\Delta^{17}\text{O-NO}_3^-$ values were substituted into equation (3), mean proportions of atmospheric NO_3^- were 54, 9, 10, and 7% in the Stage 0, 1, 2, and 3 watersheds, respectively. The $\Delta^{17}\text{O}$ -based estimate for the Stage 0 watershed (54%) should be interpreted with caution, as samples collected on only four dates during 2010 had adequate NO_3^- -N concentrations for $\Delta^{17}\text{O-NO}_3^-$ analysis (for comparison, mean proportions of atmospheric NO_3^- in the Stage 1, 2, and 3 watersheds calculated for these four sampling dates were 20, 11, and 10%, respectively). This caveat notwithstanding, both $\delta^{18}\text{O}$ - and $\Delta^{17}\text{O}$ -based estimates indicate that proportions of atmospheric

Table 3. Weighted Means (SE) of Growing Season and Dormant Season NO_3^- -N Concentrations and Isotope Values

N Sat. Stage	$[\text{NO}_3^- \text{-N}]$ (mg L^{-1})		$\delta^{15}\text{N-NO}_3^-$ (‰)		$\delta^{18}\text{O-NO}_3^-$ (‰)		$\Delta^{17}\text{O-NO}_3^-$ (‰)	
	Growing	Dormant	Growing	Dormant	Growing	Dormant	Growing	Dormant
0	0.00 (0.00)	0.00 (0.00)	-0.6 (1.2)	+2.0 (0.9)	+10.3 ^a (7.7)	+58.3 ^b (5.9)	+3.3 (1.7)	+23.9 <i>na</i>
1	0.20 ^a (0.02)	0.26 ^b (0.01)	+3.1 ^a (0.2)	+1.8 ^b (0.1)	-5.1 ^a (1.2)	-0.5 ^b (0.7)	-0.6 ^a (0.4)	+0.7 ^b (0.3)
2	0.35 ^a (0.04)	0.50 ^b (0.02)	+2.7 ^a (0.1)	+1.9 ^b (0.1)	-4.0 ^a (1.0)	+0.1 ^b (0.5)	-0.1 ^a (0.3)	+0.8 ^b (0.2)
3	0.71 ^a (0.03)	0.87 ^b (0.02)	+3.9 ^a (0.2)	+2.3 ^b (0.2)	-4.8 ^a (0.6)	-1.6 ^b (0.4)	-0.8 (0.2)	+0.1 (0.1)
Precip	0.15 (0.03)	0.21 (0.03)	-0.7 (0.6)	+0.6 (0.6)	+62.5 ^a (1.4)	+68.0 ^b (1.4)	+19.4 ^a (0.9)	+23.6 ^b (1.0)

Growing and dormant season means followed by different letters are significantly different at $\alpha = 0.05$. *na* denotes that the standard error could not be calculated due to $n = 1$.

NO_3^- in streams were much lower in the Stage 1, 2, and 3 (hardwood) watersheds than in the Stage 0 (conifer) watershed, and mean proportions of atmospheric NO_3^- tended to decrease with increasing mean stream NO_3^- -N concentrations. For all watersheds, the mean proportions of atmospheric NO_3^- in streams estimated by $\delta^{18}\text{O-NO}_3^-$ were greater than those based on $\Delta^{17}\text{O-NO}_3^-$.

In contrast to the pattern of atmospheric NO_3^- proportions in streams, total fluxes of atmospheric NO_3^- -N in streams increased with increasing mean NO_3^- -N concentration (Table 4). During the study period, total stream fluxes of atmospheric NO_3^- -N were 2.9, 117.8, 207.5, and 338.5 g ha^{-1} in the Stage 0, 1, 2, and 3 watersheds, respectively (Table 4). These $\Delta^{17}\text{O}$ -based flux estimates do not account for contributions of atmospheric NH_4^+ that may have been oxidized to NO_3^- prior to export from the watersheds. Therefore, these estimates represent only that fraction of total stream N export contributed directly from atmospheric NO_3^- . Total atmospheric NO_3^- -N deposition (wet + dry) at Fernow during 2010 was 3.28 kg ha^{-1} and total inorganic N deposition (wet + dry NO_3^- -N and NH_4^+ -N) was 5.25 kg ha^{-1} (U.S. Environmental Protection Agency 2014).

DISCUSSION

Biological and Hydrologic Influences on Stream Nitrate Sources and Export

The patterns of atmospheric NO_3^- export from the study watersheds at Fernow contradict our initial hypothesis of increasing proportions of atmospheric NO_3^- in streams with increasing N saturation status. However, on a flux basis, atmospheric NO_3^- contributions to streams increased with increasing N saturation stage, consistent with the results of Durka and others (1994) who observed the greatest

atmospheric NO_3^- fluxes from forests in a deposition-induced state of decline. Our results also demonstrate a comparatively stronger response of microbial NO_3^- fluxes to increasing N saturation stage relative to atmospheric NO_3^- fluxes, suggesting that while the capacity for biological N cycling may not decrease with progressive N saturation, the capacity for N retention within watersheds can diminish.

The study watersheds at Fernow reflect important differences in biological and hydrologic ecosystem attributes. Perhaps the most obvious difference is that of the species composition in the Stage 0 watershed, the Norway spruce monoculture. Many previous studies have demonstrated the influence of species composition on soil characteristics related to N cycling and export (Buldgen and others 1983; Lovett and others 2002, 2004; Templer and others 2005; Kelly and others 2011; Prescott and Grayston 2013). The high $\Delta^{17}\text{O-NO}_3^-$ values in the Stage 0 watershed indicate that atmospheric deposition is a more important source of stream NO_3^- than microbial nitrification in this conifer-dominated watershed. Although it is possible that the low stream NO_3^- -N concentrations and limited $\Delta^{17}\text{O-NO}_3^-$ data in the Stage 0 watershed resulted in a biased estimate of atmospheric NO_3^- contributions in this stream, additional data support the observation that microbial nitrification constitutes a smaller proportion of total stream NO_3^- in the Stage 0 watershed than in the Stage 1, 2, and 3 watersheds. First, although $\delta^{18}\text{O}$ is a less conservative tracer of atmospheric NO_3^- than $\Delta^{17}\text{O}$ (Michalski and others 2004; Kendall and others 2007; Riha and others 2014), the larger dataset ($n = 12$) and high isotope values of $\delta^{18}\text{O-NO}_3^-$ in the Stage 0 watershed (median = +41.7‰) support the conclusion that atmospheric deposition is a major source of stream NO_3^- . Second, the mean A horizon C:N in the

Table 4. Characteristics of Annual Stream Discharge, NO_3^- -N Concentrations, Stream NO_3^- -N Fluxes, and N Retention in the Study Watersheds

N sat. stage	Total discharge (mm)	Discharge-weighted mean stream $[\text{NO}_3^-]$ (mg L^{-1})	Total stream NO_3^- -N flux (g ha^{-1})	Total atmospheric NO_3^- -N flux in streams, based on $\Delta^{17}\text{O}$ (g ha^{-1}) ¹	Discharge-weighted mean proportion of atmospheric NO_3^- in streams, based on $\Delta^{17}\text{O}$ (%) ¹	Discharge-weighted mean proportion of atmospheric NO_3^- in streams, based on $\delta^{18}\text{O}$ (%)	Watershed N retention, based on atmospheric NO_3^- inputs (%) ²	Watershed N retention, based on atmospheric $\text{NO}_3^- + \text{NH}_4^+$ inputs (%) ²
0	199.2	0.00	5.4	2.9	54 ($n = 4$)	76 ($n = 12$)	99	99
1	567.1	0.25	1308.8	117.8	9 ($n = 38$)	14 ($n = 38$)	60	75
2	449.8	0.48	2075.2	207.5	10 ($n = 32$)	12 ($n = 32$)	37	60
3	577.7	0.84	4835.7	338.5	7 ($n = 41$)	9 ($n = 41$)	-47	8

¹ Values for the Stage 0 watershed should be interpreted with caution, as they were calculated using only four sampling dates during 2010 for which $\Delta^{17}\text{O}$ - NO_3^- data were available (for comparison, total atmospheric NO_3^- -N fluxes in the Stage 1, 2, and 3 watersheds for these four dates were 7.9, 63, and 16.7 g ha^{-1} , respectively).

² Watershed N retention estimates were calculated using total (wet + dry) deposition of NO_3^- -N or inorganic N (NO_3^- -N and NH_4^+ -N) measured at the National Atmospheric Deposition Program and EPA Clean Air Status and Trends Network sites at Fernow during 2010. Total atmospheric NO_3^- -N deposition was 3.28 kg ha^{-1} and total inorganic N deposition was 5.25 kg ha^{-1} .

Stage 0 watershed is higher than that in the Stage 3 watershed (18 vs. 15 in the Stage 0 and 3 watersheds, respectively; (Kelly and others 2011)), and the Stage 2 watershed (mean C:N = 16; Christ and others 2002). It is possible that A horizon C:N is more spatially variable in the Stage 2 and 3 watersheds due to greater species diversity in these mixed hardwood-dominated watersheds compared to the Norway spruce-dominated Stage 0 watershed, potentially influencing mineralization and nitrification rates (Lovett and others 2004). Typically, litter from conifer species is more recalcitrant (that is, higher C:N and lignin:N) than litter from hardwood species such as maple and tulip poplar (Lovett and others 2004; Kelly 2010), and previous studies have demonstrated significant negative correlations between soil C:N and net nitrification rates (Christ and others 2002; Ross and others 2004; Christenson and others 2009; Kelly and others 2011). In the Stage 0 watershed, the mean annual net nitrification rate of $4.2 \text{ kg N ha}^{-1} \text{ y}^{-1}$ (Kelly 2010) is 27 times lower than the rates measured by Gilliam and others (2001) in the Stage 2 and 3 watersheds (114 and $115 \text{ kg N ha}^{-1} \text{ y}^{-1}$ respectively). This markedly lower rate of microbial NO_3^- production in the Stage 0 watershed contributes to both the low proportions of microbial NO_3^- in the stream and the low total NO_3^- -N flux from this watershed. The stream NO_3^- source dynamics observed in the study watersheds have important implications for our understanding of the ecosystem effects of N deposition. For example, the higher proportions of atmospheric NO_3^- in the Stage 0 stream suggest that this watershed might show the strongest immediate response to changing rates of atmospheric N deposition. In contrast, the overwhelming contributions of microbial NO_3^- to the Stage 1, 2, and 3 streams suggest that a reduction in rates of atmospheric N deposition to these watersheds might yield only minor immediate reductions in stream NO_3^- concentrations and fluxes. In the longer term, reductions in N deposition would likely reduce the level of N cycling and nitrification in these watersheds, resulting in a decline in the microbial fraction of exported NO_3^- .

Although the relative contributions of NO_3^- from atmospheric and microbial sources influence stream NO_3^- -N concentrations and fluxes, watershed hydrologic status is an additional key driver of microbial and atmospheric NO_3^- export to streams (Creed and Band 1998; Adams and others 2014; Sebastyen and others 2014). In all of the study watersheds, $\delta^{18}\text{O}$ - NO_3^- and $\Delta^{17}\text{O}$ - NO_3^- values were generally higher in stream samples collected during stormflow, and both $\delta^{18}\text{O}$ - NO_3^- and $\Delta^{17}\text{O}$ -

NO_3^- were positively correlated with discharge ($P < 0.05$) (except $\Delta^{17}\text{O}-\text{NO}_3^-$ in the Stage 0 watershed, due to the limited $\Delta^{17}\text{O}-\text{NO}_3^-$ dataset in this watershed). In the Stage 1, 2, and 3 watersheds, 54 to 61% of total NO_3^- export and 62 to 77% of total atmospheric NO_3^- export during 2010 occurred on nine sampling dates, coincident with stormflow. This suggests that most of the total NO_3^- (and atmospheric NO_3^- in particular) that was exported from these watersheds during 2010 occurred during stormflow. The role of hydrologic drivers in regulating atmospheric NO_3^- export is further demonstrated by the high $\Delta^{17}\text{O}-\text{NO}_3^-$ values measured in all watersheds on 23 February, coincident with the rising limb of a snowmelt event. Using the highest precipitation $\Delta^{17}\text{O}-\text{NO}_3^-$ value measured during the study period (+29.1‰) and the lowest $\Delta^{17}\text{O}-\text{NO}_3^-$ value measured in each watershed (and 0‰ for the Stage 0 watershed) for the nitrification end-member in equation (3), conservative estimates of atmospheric NO_3^- in streams were 82, 22, 15, and 15% of total stream NO_3^- in the Stage 0, 1, 2, and 3 watersheds, respectively, on 23 February. Rapid routing of melt water to streams and decreased biological activity during the winter could have facilitated elevated proportions of atmospheric NO_3^- in streams on this date. Several studies in forested watersheds have also reported elevated atmospheric NO_3^- contributions to streams during snowmelt (Burns and Kendall 2002; Sebastyen and others 2008; Goodale and others 2009; Pellerin and others 2012). That the largest proportions of atmospheric NO_3^- in all streams occurred on the same date suggests that a common factor (for example, hydrologic status) influenced atmospheric NO_3^- export from the study watersheds during this stormflow event. However, when examined over the entire study period, differing biogeochemical responses among some of the study watersheds indicate that hydrologic and biological drivers may interact to regulate NO_3^- export dynamics over longer temporal scales.

Although the lowest mean proportion of atmospheric NO_3^- (7%) occurred in the Stage 3 watershed, total NO_3^- -N concentrations and discharge were highest in this watershed resulting in the greatest fluxes of both total and atmospheric NO_3^- -N (Table 4). In contrast, the Stage 1 watershed had a slightly greater mean proportion of atmospheric NO_3^- (9%) and nearly identical total annual discharge to that in the Stage 3 watershed, but total and atmospheric NO_3^- -N fluxes in the Stage 1 watershed were around 3 times less than those in the Stage 3 watershed. Thus, despite identical atmospheric deposition inputs and nearly identical

patterns of discharge during the study period, these two watersheds exhibited differing total and atmospheric NO_3^- exports during 2010. The difference in NO_3^- -N fluxes indicates that retention of atmospheric NO_3^- in the Stage 1 watershed is greater than that in the Stage 3 watershed. This may be due to biological differences (for example, degree of NO_3^- retention in vegetation and soils, possibly due to lower nitrification rates in the Stage 1 watershed), hydrologic differences (for example, hydrologic transit times along flowpaths), or some combination of these (and possibly additional) factors. Although the predominant hydrologic source areas and flowpaths have not been characterized in these watersheds, we might expect the steeper slopes in the Stage 1 watershed (mean = 30.6% versus 24.8% in the Stage 3 watershed) to facilitate equal or greater fluxes of atmospheric NO_3^- to streams if hydrology were the primary driver of atmospheric NO_3^- export. That less atmospheric NO_3^- was exported from the Stage 1 watershed compared to the Stage 3 watershed despite similarities in discharge and atmospheric inputs suggests that the capacity for retention of atmospheric NO_3^- may be lower in the Stage 3 watershed. Indeed, the Stage 3 watershed served as a net source of NO_3^- -N relative to the amount of atmospheric NO_3^- -N deposited during 2010 (Table 4). However, quantification of the relative strengths of vegetation and soil N sinks in the two watersheds would be required to validate this speculation.

Although the higher proportions of atmospheric NO_3^- in streams during stormflow at Fernow are in agreement with the results of other studies that have examined atmospheric NO_3^- dynamics during hydrologic events (Burns and Kendall 2002; Sebastyen and others 2008, 2014; Pellerin and others 2012; Riha and others 2014), it is important to reiterate that the proportions of atmospheric NO_3^- we observed in streams are generally small throughout the year (range of median atmospheric NO_3^- proportions in all streams = 3 to 39%). Indeed, much larger discharges than those observed during the snowmelt event on 23 February were recorded from all of the watersheds during 2010 but these events occurred during the growing season, resulting in only marginal increases in the proportion of atmospheric NO_3^- in streams. Therefore, although hydrologic status is an important driver of atmospheric NO_3^- dynamics in these streams, extensive microbial NO_3^- production—particularly in the Stage 2 and 3 watersheds (Gilliam and others 2001)—also influences NO_3^- source dynamics and total NO_3^- -N flux.

Seasonal Nitrate Isotope Dynamics

Temporal patterns of triple NO_3^- isotopes provide insight into the biological N cycling dynamics of the study watersheds. A strongly seasonal pattern of stream $\delta^{15}\text{N}-\text{NO}_3^-$ values occurred in the Stage 1, 2, and 3 watersheds (Figure 3). The sharp increase in $\delta^{15}\text{N}-\text{NO}_3^-$ values from early April through late August may have resulted from mass-dependent fractionations during nitrification and/or plant uptake. This increase was likely not due to denitrification, as a concomitant increase in $\delta^{18}\text{O}-\text{NO}_3^-$ values (characteristic of denitrification activity; Kendall 1998) did not occur in any of the streams. Although mineralization of organic N to NH_4^+ does not result in large isotopic fractionations (Kendall 1998), oxidation of NH_4^+ to NO_3^- can yield fractionations ranging from -12‰ to -29‰ (Shearer and Kohl 1986). Thus, as the ratio of net nitrification to net mineralization increases, the $\delta^{15}\text{N}-\text{NO}_3^-$ in soils should approach that of the original soil NH_4^+ pool (Spoelstra and others 2007). Net nitrification rates in the Stage 2 and 3 watersheds are high during the growing season—reaching nearly 100% of net N mineralization rates—and soil N pools remain well above $0\text{ g NO}_3^- \text{ N m}^{-2}$ throughout the year, indicating that microbial $\text{NO}_3^- \text{ N}$ production exceeds plant uptake in these watersheds (Gilliam and others 2001). Preferential uptake of the lighter ^{14}N isotope during the growing season may have yielded an isotopically-enriched residual soil NO_3^- pool (Högberg 1997; Kendall and others 2007; Templer and others 2007) from which stream NO_3^- was derived. Burns and Kendall (2002) reported $\delta^{15}\text{N}-\text{NO}_3^-$ values ranging from $+9$ to $+16\text{‰}$ in O horizon soils incubated at 22°C compared to $+1.5\text{‰}$ in soils incubated at 4°C . The authors suggested that greater cycling of N through mineralization, nitrification, and immobilization processes may have yielded higher isotope values in the soils incubated at higher temperatures. Similar processes could have contributed to the enrichment in $\delta^{15}\text{N}$ values observed during the growing season in the Stage 1, 2, and 3 watersheds at Fernow. Although plant N demand decreases sharply at the end of the growing season, soil microbial processes such as nitrification can continue throughout the dormant season under an insulating snowpack (Campbell and others 2005). Decreased plant N demand at the end of the growing season and throughout the dormant season may similarly explain the lower $\delta^{15}\text{N}-\text{NO}_3^-$ values observed in streams during this time, as isotopic fractionation in the residual soil NO_3^- pool would have been less pronounced. The low $\delta^{15}\text{N}-$

NO_3^- values observed in the Stage 0 watershed (median = $+0.7\text{‰}$), their similarity to precipitation $\delta^{15}\text{N}-\text{NO}_3^-$ values (median = $+0.3\text{‰}$), and the absence of seasonal isotopic enrichment in this watershed suggest that biologically mediated isotopic fractionations in the soil NO_3^- pool were less extensive in this watershed. This is in agreement with previous studies at Fernow that reported much lower rates of net nitrification in the Stage 0 watershed compared to the Stage 2 and 3 watersheds (Gilliam and others 2001; Kelly 2010).

In contrast to the seasonal enrichment of stream $\delta^{15}\text{N}-\text{NO}_3^-$ values in the Stage 1, 2, and 3 watersheds, stream $\delta^{18}\text{O}-\text{NO}_3^-$ values were generally lower during the growing season in all watersheds (Figure 3; Table 3). The absence of a seasonal enrichment in stream $\delta^{18}\text{O}-\text{NO}_3^-$ values similar to that observed for $\delta^{15}\text{N}-\text{NO}_3^-$ is unexpected, as both are subject to mass-dependent fractionation during biological processes and exhibit similar isotope effects during biological assimilation (Granger and others 2004; Deutsch and others 2009). It is possible that a seasonal enrichment in $\delta^{18}\text{O}-\text{NO}_3^-$ values did occur similar to that observed in $\delta^{15}\text{N}-\text{NO}_3^-$ values, but the effect was obscured by the concomitant decline in proportions of atmospheric NO_3^- in streams during the growing season. Due to the greater disparity in $\delta^{18}\text{O}-\text{NO}_3^-$ values between atmospheric and microbial sources relative to the differences in $\delta^{15}\text{N}-\text{NO}_3^-$ values, it is possible that decreasing proportions of atmospheric NO_3^- in streams obscured any uptake-induced enrichment that may have occurred in $\delta^{18}\text{O}-\text{NO}_3^-$ values. In addition, Burns and Kendall (2002) attributed greater variation in $\delta^{15}\text{N}$ values compared to $\delta^{18}\text{O}$ values of NO_3^- in a soil incubation experiment to differences in reaction pathways of nitrogen and oxygen during microbial NO_3^- processing. Whereas the original isotopic signature of N is partially preserved during N cycling processes, the isotopic composition of oxygen is reset with each nitrification cycle (Burns and Kendall 2002). The oxygen isotopic composition of microbial NO_3^- is reset through variable contributions of oxygen atoms from soil water and O_2 in soils during nitrification (Buchwald and Casciotti 2010; Snider and others 2010); such exchange processes may have also contributed to the divergent seasonal patterns in $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ of NO_3^- observed in the study watersheds.

In contrast to the seasonal pattern in $\delta^{18}\text{O}-\text{NO}_3^-$ values, $\Delta^{17}\text{O}-\text{NO}_3^-$ values in weekly stream samples from the Stage 1, 2, and 3 watersheds were less variable from January to June (Figure 3), although significant seasonal differences were observed in the

Stage 1 and 2 watersheds ($P < 0.05$; Table 3). Consequently, estimates of atmospheric NO_3^- proportions in streams calculated using $\delta^{18}\text{O}\text{-NO}_3^-$ were higher during the dormant season than proportions calculated using $\Delta^{17}\text{O}\text{-NO}_3^-$ values (Figure 4). The




Figure 4. Estimated percent atmospheric NO_3^- in weekly stream samples in hardwood watersheds at Fernow using $\Delta^{17}\text{O}$ and $\delta^{18}\text{O}$ of NO_3^- values in a two end-member isotope mixing model. $\delta^{18}\text{O}$ -based percentages over-estimate atmospheric NO_3^- contributions to streams during the dormant season (shown by circles), whereas percent estimates in growing season samples (shown by crosses) plot closer to the 1:1 line (shown by the dotted line) in all watersheds.

greatest differences between $\delta^{18}\text{O}$ - and $\Delta^{17}\text{O}$ -based estimates occurred in January and February, when $\delta^{18}\text{O}\text{-NO}_3^-$ values were highest in streams; $\delta^{18}\text{O}$ -based estimates were up to 12% higher than $\Delta^{17}\text{O}$ -based estimates during this time. This may be attributable to seasonal differences in the $\delta^{18}\text{O}$ values of water used by soil microbes during nitrification, as previously discussed. If seasonal variations in the oxygen isotopic composition of atmospheric NO_3^- alone were responsible for the higher dormant season $\delta^{18}\text{O}\text{-NO}_3^-$ values observed in streams, this would have also been reflected in the pattern of $\Delta^{17}\text{O}\text{-NO}_3^-$ values of stream samples in all of the watersheds. That the $\delta^{18}\text{O}$ -based estimates of atmospheric NO_3^- in streams are greater than those based on $\Delta^{17}\text{O}\text{-NO}_3^-$ and show a more distinctly seasonal pattern suggests that biologically mediated isotope fractionations resulted in the biased $\delta^{18}\text{O}$ -based estimates of atmospheric NO_3^- . Previous studies have also attributed higher $\delta^{18}\text{O}$ -based estimates of atmospheric NO_3^- in streams (compared to $\Delta^{17}\text{O}\text{-NO}_3^-$ -based estimates) to the influence of microbial activity (Riha and others 2014). These results demonstrate the advantage of using $\Delta^{17}\text{O}\text{-NO}_3^-$ over $\delta^{18}\text{O}\text{-NO}_3^-$ for more precise apportionment of atmospheric and biological NO_3^- contributions to streams.

CONCLUSIONS

Four nearly adjacent watersheds receiving identical atmospheric NO_3^- inputs but differing significantly in long-term patterns of stream NO_3^- export showed variable $\delta^{15}\text{N}$, $\delta^{18}\text{O}$, and $\Delta^{17}\text{O}$ dynamics of stream NO_3^- during 2010. Our results indicate that:

- (1) Atmospheric deposition can serve as the dominant source of NO_3^- in some streams, particularly in watersheds characterized by low microbial nitrification rates and low stream NO_3^- concentrations.
- (2) Both flux-based and concentration-based estimates should be considered when evaluating NO_3^- source contributions to streams, as the relative importance of atmospheric NO_3^- may differ depending on which approach is used.
- (3) Watershed hydrologic status influences the export of atmospheric NO_3^- to streams, with greater proportions of atmospheric NO_3^- in stormflow than baseflow.
- (4) Nitrate source apportionments based on $\delta^{18}\text{O}\text{-NO}_3^-$ can over-estimate contributions from atmospheric deposition relative to $\Delta^{17}\text{O}\text{-NO}_3^-$ -based estimates. Because it is not affected by mass-dependent fractionations that commonly

occur during biological processes, $\Delta^{17}\text{O}-\text{NO}_3^-$ is a more robust and conservative tracer of atmospheric NO_3^- than $\delta^{18}\text{O}-\text{NO}_3^-$.

ACKNOWLEDGMENTS

This work was supported by NSF Grant #0910521 (to EME) and Agriculture and Food Research Initiative Grant #2012-67011-19663 (to LAR) from the USDA National Institute of Food and Agriculture. We thank Christopher Cassidy for assistance with sample collection and processing, Katherine Redling for isotopic analysis, and the helpful comments of three anonymous reviewers during the preparation of this manuscript.

REFERENCES

- Aber JD, Nadelhoffer KJ, Steudler P, Melillo JM. 1989. Nitrogen saturation in northern forest ecosystems. *Bioscience* 39:378–86.
- Aber J, Goodale CL, Ollinger S, Smith ML, Magill A, Martin ME, Hallet RA, Stoddard JL. 2003. Is nitrogen deposition altering the nitrogen status of northeastern forests? *Bioscience* 53:375–89.
- Aber J, McDowell W, Nadelhoffer K, Magill A, Berntson G, Kamakea M, McNulty S, Currie W, Rustad L, Fernandez I. 1998. Nitrogen saturation in temperate forest ecosystems. *Bioscience* 48:921–34.
- Adams MB, DeWalle DR, Peterjohn WT, Gilliam FS, Sharpe WE, Williard KW. 2006. Soil chemical response to experimental acidification treatments. In: *The Fernow Watershed Acidification Study*. Springer. pp 41–69.
- Adams MB, Edwards PJ, Kochenderfer JN, Wood F. 2003. Fifty years of watershed research on the Fernow Experimental Forest, WV: effects of forest management and air pollution in hardwood forests. First interagency conference on research in the watersheds. Benson: USDA-ARS. p 391–6.
- Adams MB, Knoepp JD, Webster JR. 2014. Inorganic nitrogen retention by watersheds at Fernow Experimental Forest and Coweeta Hydrologic Laboratory. *Soil Sci Soc Am J* 78:S84–94.
- Adams MB, Kochenderfer JN, Edwards PJ. 2007. The Fernow watershed acidification study: ecosystem acidification, nitrogen saturation and base cation leaching. *Water Air Soil Pollut Focus* 7:267–73.
- Barnes RT, Raymond PA, Casciotti KL. 2008. Dual isotope analyses indicate efficient processing of atmospheric nitrate by forested watersheds in the northeastern US. *Biogeochemistry* 90:15–27.
- Buchwald C, Casciotti KL. 2010. Oxygen isotopic fractionation and exchange during bacterial nitrite oxidation. *Limnol Oceanogr* 55:1064–74.
- Buldgen P, Dubois D, Remacle J. 1983. Principal component analysis applied to nutrient balances in organic layers of beech and spruce forests. *Soil Biol Biochem* 15:511–18.
- Burns DA, Kendall C. 2002. Analysis of $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ to differentiate NO_3^- sources in runoff at two watersheds in the Catskill Mountains of New York. *Water Resour Res* 38:9–1.
- Campbell JL, Mitchell MJ, Groffman PM, Christenson LM, Hardy JP. 2005. Winter in northeastern North America: a critical period for ecological processes. *Front Ecol Environ* 3:314–22.
- Casciotti KL, Sigman DM, Hastings MG, Böhlke JK, Hilkert A. 2002. Measurement of the oxygen isotopic composition of nitrate in seawater and freshwater using the denitrifier method. *Anal Chem* 74:4905–12.
- Christenson LM, Lovett GM, Weathers KC, Arthur MA. 2009. The influence of tree species, nitrogen fertilization, and soil C to N ratio on gross soil nitrogen transformations. *Soil Sci Soc Am J* 73:638–46.
- Christ MJ, Peterjohn WT, Cumming JR, Adams MB. 2002. Nitrification potentials and landscape, soil and vegetation characteristics in two central Appalachian watersheds differing in NO_3^- export. *For Ecol Manage* 159:145–58.
- Coplen TB, Böhlke JK, Casciotti KL. 2004. Using dual-bacterial denitrification to improve $\delta^{15}\text{N}$ determinations of nitrates containing mass-independent ^{17}O . *Rapid Commun Mass Spectrom* 18:245–50.
- Creed IF, Band LE. 1998. Export of nitrogen from catchments within a temperate forest: evidence for a unifying mechanism regulated by variable source area dynamics. *Water Resour Res* 34:3105–20.
- Deutsch B, Voss M, Fischer H. 2009. Nitrogen transformation processes in the Elbe River: distinguishing between assimilation and denitrification by means of stable isotope ratios in nitrate. *Aquat Sci* 71:228–37.
- Durka W, Schulze ED, Gebauer G, Voerkellust S. 1994. Effects of forest decline on uptake and leaching of deposited nitrate determined from ^{15}N and ^{18}O measurements. *Nature* 372:22.
- Fernandez IJ, Adams MB, SanClements MD, Norton SA. 2010. Comparing decadal responses of whole-watershed manipulations at the Bear Brook and Fernow experiments. *Environ Monit Assess* 171:149–61.
- Galloway JN, Aber JD, Erisman JW, Seitzinger SP, Howarth RW, Cowling EB, Cosby BJ. 2003. The nitrogen cascade. *Bioscience* 53:341–56.
- Gilliam FS, Yurish BM, Adams MB. 2001. Temporal and spatial variation of nitrogen transformations in nitrogen-saturated soils of a central Appalachian hardwood forest. *Can J For Res* 31:1768–85.
- Goodale CL, Thomas SA, Fredriksen G, Elliott EM, Flinn KM, Butler TJ, Walter MT. 2009. Unusual seasonal patterns and inferred processes of nitrogen retention in forested headwaters of the Upper Susquehanna River. *Biogeochemistry* 93:197–218.
- Granger J, Sigman DM, Needoba JA, Harrison PJ. 2004. Coupled nitrogen and oxygen isotope fractionation of nitrate during assimilation by cultures of marine phytoplankton. *Limnol Oceanogr* 49:1763–73.
- Högberg P. 1997. Tansley Review No. 95 ^{15}N natural abundance in soil-plant systems. *New Phytol* 137:179–203.
- Kaiser J, Hastings MG, Houlton BZ, Röckmann T, Sigman DM. 2007. Triple oxygen isotope analysis of nitrate using the denitrifier method and thermal decomposition of N_2O . *Anal Chem* 79:599–607.
- Kelly CN. 2010. Carbon and nitrogen cycling in watersheds of contrasting vegetation types in the Fernow Experimental Forest, West Virginia. Doctoral dissertation, Virginia Polytechnic Institute and State University.
- Kelly C, Schoenholtz S, Adams M. 2011. Soil properties associated with net nitrification following watershed conversion

- from Appalachian hardwoods to Norway spruce. *Plant Soil* 344:361–76.
- Kendall C. 1998. Tracing nitrogen sources and cycling in catchments. In: Kendall C, McDonnell JJ, Eds. *Isotope Tracers in Catchment Hydrology*. Amsterdam: Elsevier. p 519–76.
- Kendall C, Elliott EM, Wankel SD. 2007. Tracing anthropogenic inputs of nitrogen to ecosystems. In: Michener R, Lajtha K, Eds. *Stable Isotopes in Ecology and Environmental Science*, Vol. 2. Oxford: Blackwell Publishing. p 375–449.
- Kochenderfer JN. 2007. Fernow and the Appalachian hardwood region. In: Adams MB, DeWalle DR, Hom JR, Eds. *The Fernow Watershed Acidification Study*. Dordrecht, The Netherlands: Springer. p 17–39.
- Lovett G, Goodale C. 2011. A new conceptual model of nitrogen saturation based on experimental nitrogen addition to an oak forest. *Ecosystems* 14:615–31.
- Lovett GM, Weathers KC, Arthur MA. 2002. Control of nitrogen loss from forested watersheds by soil carbon: nitrogen ratio and tree species composition. *Ecosystems* 5:0712–18.
- Lovett GM, Weathers KC, Arthur MA, Schultz JC. 2004. Nitrogen cycling in a northern hardwood forest: do species matter? *Biogeochemistry* 67:289–308.
- Michalski G, Meixner T, Fenn M, Hernandez L, Sirulnik A, Allen E, Thiemens M. 2004. Tracing atmospheric nitrate deposition in a complex semiarid ecosystem using $\Delta^{17}\text{O}$. *Environ Sci Technol* 38:2175–81.
- Pardo LH, Kendall C, Pett-Ridge J, Chang CCY. 2004. Evaluating the source of streamwater nitrate using $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ in nitrate in two watersheds in New Hampshire, USA. *Hydrol Process* 18:2699–712.
- Pellerin BA, Saraceno JF, Shanley JB, Sebestyen SD, Aiken GR, Wollheim WM, Bergamaschi BA. 2012. Taking the pulse of snowmelt: in situ sensors reveal seasonal, event and diurnal patterns of nitrate and dissolved organic matter variability in an upland forest stream. *Biogeochemistry* 108:183–98.
- Peterjohn WT, Adams MB, Gilliam FS. 1996. Symptoms of nitrogen saturation in two central Appalachian hardwood forest ecosystems. *Biogeochemistry* 35:507–22.
- Prescott CE, Grayston SJ. 2013. Tree species influence on microbial communities in litter and soil: current knowledge and research needs. *For Ecol Manage* 309:19–27.
- Reinhart KG, Eschner A, Trimble GR. 1963. Effect on streamflow of four forest practices in the mountains of West Virginia. Research Paper NE-1. Upper Darby, PA: US Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 79 p.
- Riha K, Michalski G, Gallo E, Lohse K, Brooks P, Meixner T. 2014. High atmospheric nitrate inputs and nitrogen turnover in semi-arid urban catchments. *Ecosystems* 1–17.
- Ross DS, Lawrence GB, Fredriksen G. 2004. Mineralization and nitrification patterns at eight northeastern USA forested research sites. *For Ecol Manage* 188:317–35.
- SAS Institute, Inc. 2011. *SAS/STAT user's guide: Statistics*, Release 9.3 edition. Cary, NC.: SAS Institute, Inc.
- Sebestyen SD, Boyer EW, Shanley JB, Kendall C, Doctor DH, Aiken GR, Ohte N. 2008. Sources, transformations, and hydrological processes that control stream nitrate and dissolved organic matter concentrations during snowmelt in an upland forest. *Water Resour Res* 44:W12410.
- Sebestyen SD, Shanley JB, Boyer EW, Kendall C, Doctor DH. 2014. Coupled hydrological and biogeochemical processes controlling variability of nitrogen species in streamflow during autumn in an upland forest. *Water Resour Res* 50:1569–91.
- Shearer G, Kohl DH. 1986. N_2 -fixation in field settings: estimations based on natural ^{15}N abundance. *Funct Plant Biol* 13:699–756.
- Sigman DM, Casciotti KL, Andreani M, Barford C, Galanter M, Böhlke JK. 2001. A bacterial method for the nitrogen isotopic analysis of nitrate in seawater and freshwater. *Anal Chem* 73:4145–53.
- Snider DM, Spoelstra J, Schiff SL, Venkiteswaran JJ. 2010. Stable oxygen isotope ratios of nitrate produced from nitrification: ^{18}O -labeled water incubations of agricultural and temperate forest soils. *Environ Sci Technol* 44:5358–64.
- Spoelstra J, Schiff SL, Hazlett PW, Jeffries DS, Semkin RG. 2007. The isotopic composition of nitrate produced from nitrification in a hardwood forest floor. *Geochim Cosmochim Acta* 71:3757–71.
- Stoddard JL. 1994. Long-term changes in watershed retention of nitrogen. Its causes and aquatic consequences. In: Baker L, Ed. *Environmental Chemistry of Lakes and Reservoirs*. Washington, DC: American Chemical Society. p 223–84.
- Templer PH, Arthur MA, Lovett GM, Weathers KC. 2007. Plant and soil natural abundance $\delta^{15}\text{N}$: indicators of relative rates of nitrogen cycling in temperate forest ecosystems. *Oecologia* 153:399–406.
- Templer PH, Lovett GM, Weathers KC, Findlay SE, Dawson TE. 2005. Influence of tree species on forest nitrogen retention in the Catskill Mountains, New York, USA. *Ecosystems* 8:1–16.
- Tiner RW. 1993. The primary indicators method—a practical approach to wetland recognition and delineation in the United States. *Wetlands* 13:50–64.
- U.S. Environmental Protection Agency. 2014. U.S. Environmental Protection Agency Clean Air Markets Division Clean Air Status and Trends Network (CASTNET). www.epa.gov/castnet. Last accessed 21/10/2014.
- Williard KW, DeWalle DR, Edwards PJ, Sharpe WE. 2001. ^{18}O isotopic separation of stream nitrate sources in mid-Appalachian forested watersheds. *J. Hydrol* 252:174–88.