

THE PROBABILITY DISTRIBUTION OF BINARY PULSAR COALESCENCE RATES. II. NEUTRON STAR–WHITE DWARF BINARIES

CHUNGLEE KIM,¹ VASSILIKI KALOGERA,¹ DUNCAN R. LORIMER,² AND TIFFANY WHITE¹

Received 2004 February 5; accepted 2004 August 7

ABSTRACT

We consider the statistics of pulsar binaries with white dwarf companions (NS-WD). Using the statistical analysis method developed by Kim and coworkers, we calculate the Galactic coalescence rate of NS-WD binaries due to gravitational wave emission. We find that the most likely values for the total Galactic coalescence rate (\mathcal{R}_{tot}) of NS-WD binaries lie in the range 0.2–10 Myr⁻¹, depending on different assumed pulsar population models. For our reference model, we obtain $\mathcal{R}_{\text{tot}} = 4.11_{-2.56}^{+5.25}$ Myr⁻¹ at a 68% statistical confidence level. These rate estimates are not corrected for pulsar beaming, and as such they are found to be about a factor of 20 smaller than the Galactic coalescence rate estimates for double neutron star systems. Based on our rate estimates, we calculate the gravitational wave background due to coalescing NS-WD binaries out to extragalactic distances within the frequency band of the *Laser Interferometer Space Antenna*. We find the contribution from NS-WD binaries to the gravitational wave background to be negligible.

Subject headings: binaries: close — gravitational waves — stars: neutron — white dwarfs

1. INTRODUCTION

The observed properties of double neutron star (DNS) systems, along with models of pulsar survey selection effects, have been used for many years in order to estimate the coalescence rate of DNS systems due to the emission of gravitational radiation (Narayan et al. 1991; Phinney 1991; Curran & Lorimer 1995). In Kim et al. (2003; hereafter Paper I), we presented a novel method to calculate the probability distribution of the coalescence rate estimates for pulsar binaries (see also Kalogera et al. 2004). In Paper I, we applied this method to Galactic DNS systems. Having a probability distribution at hand allowed us to calculate the most likely value for the Galactic DNS coalescence rate, as well as statistical confidence limits associated with it. These were then used to calculate the expected DNS inspiral detection rate for Laser Interferometer Gravitational-Wave Observatory (Abramovici et al. 1992). In this second paper, we extend our study to NS-WD binaries that are relevant to the future NASA/ESA mission *Laser Interferometer Space Antenna* (*LISA*; Bender et al. 1998).

There are now more than 40 neutron star–white dwarf (NS-WD) binary systems known in the Galactic disk (see, e.g., Lorimer 2001 for a review). Here, we consider the subset of NS-WD binaries that will coalesce as a result of gravitational wave (GW) emission within a Hubble time. There are currently three such coalescing binaries known: PSR J0751+1807 (Lundgren et al. 1995; Nice et al. 2004), PSR J1757–5322 (Edwards & Bailes 2001), and PSR J1141–6545 (Kaspi et al. 2000; Bailes et al. 2003). We calculate the Galactic coalescence rate of NS-WD binaries based on their observed properties using the method introduced in Paper I. The GW frequencies emitted by these systems fall within the ~0.1–100 mHz frequency band of *LISA*. Using our rate estimates, we calculate the GW amplitude due to the NS-WD binaries out to cosmological distances and compare it to the sensitivity curve of *LISA* (Larson et al. 2000), as well as the Galactic confusion noise estimates from white dwarf binaries (Hils et al. 1990; Bender & Hils 1997; Nelemans et al. 2001; Schneider et al. 2001).

The organization of the rest of this paper is as follows. In § 2, we consider the lifetimes of NS-WD binaries and summarize the techniques we use to calculate their coalescence rate. The results of these calculations are presented in § 3. In § 4 we use our rate results to calculate the expected GW background produced by NS-WD binaries. Finally, in § 5, we discuss the implications of our results.

2. COALESCING NS-WD BINARIES

In general, the coalescence rate of a binary system containing an observable radio pulsar is defined by

$$\mathcal{R} = \frac{N_{\text{PSR}}}{\tau_{\text{life}}} f_b, \quad (1)$$

where N_{PSR} is the estimated number of pulsars in our Galaxy with pulse profiles and orbital characteristics similar to those of the known systems,³ f_b is a correction factor for pulsar beaming, and τ_{life} is the lifetime of the binary system. In the following subsections, we calculate the total lifetime of a pulsar binary and derive the probability density function (PDF) of the Galactic coalescence rate, $P(\mathcal{R})$, for NS-WD binaries.

2.1. Lifetime of an NS-WD Binary

In Table 1, we summarize the observational properties and relevant lifetimes for the three pulsar systems considered in this work. We define the lifetime of a coalescing pulsar binary τ_{life} to be the sum of the current age of the observable pulsar and the remaining

¹ Department of Physics and Astronomy, Northwestern University, 2145 Sheridan Road, Evanston, IL 60208; c-kim1@northwestern.edu, vicky@northwestern.edu, tiphane3@hotmail.com.

² Jodrell Bank Observatory, University of Manchester, Macclesfield, Cheshire SK11 9DL, UK; drl@jb.man.ac.uk.

³ This is equivalent to the so-called scale factor (Narayan 1987).

TABLE 1
OBSERVATIONAL PROPERTIES OF NS-WD BINARIES

PSRs	P (ms)	\dot{P} (s s ⁻¹)	P_b (hr)	m_{wd}^a (M_\odot)	e	τ_c (Gyr)	τ_{sd} (Gyr)	τ_{mrg} (Gyr)	τ_d (Gyr)	\mathcal{N}_{PSR}	References
J0751+1807.....	3.479	8.08×10^{-21}	6.315	0.18	$<10^{-4}$	6.8	6.7	7.6	...	2900	1, 2
J1757-5322.....	8.870	2.78×10^{-20}	10.879	0.67	10^{-6}	5.1	4.9	7.8	...	1200	3
J1141-6545.....	393.9	4.29×10^{-15}	4.744	0.986	0.172	1.5×10^{-3}	...	0.6	0.104	400	4, 5

NOTES.—From left to right, the columns indicate the pulsar name, spin period P , spin-down rate \dot{P} , orbital period P_b , most probable mass of the WD companion m_{wd} , orbital eccentricity e , characteristic age τ_c , spin-down age τ_{sd} , GW merger timescale τ_{mrg} , time to reach the death line τ_d , most probable number of NS-WD systems of this type in the Galaxy, and references to this system in the literature.

^a The assumed NS mass and inclination angle i are $2.2 M_\odot$ and 78° for PSR J0751+1807 (Nice et al. 2004) and $1.35 M_\odot$ and 60° for PSR J1757-5322. For PSR J1141-6545, we adopt values given in Bailes et al. (2003).

REFERENCES.—(1) Lundgren et al. 1995; (2) Nice et al. 2004; (3) Edwards & Bailes 2001; (4) Kaspi et al. 2000; (5) Bailes et al. 2003.

lifetime of the system. If we assume that the pulsar spins down from an initial period P_0 to the currently observed value P (both in s) as a result of a nondecaying magnetic dipole radiation torque (see, e.g., Manchester & Taylor 1977), its current “spin-down” age

$$\tau_{\text{sd}} = \frac{P}{2\dot{P}} \left[1 - \left(\frac{P_0}{P} \right)^2 \right], \quad (2)$$

where \dot{P} (in s s⁻¹) is the observed period derivative. For young pulsars like PSR J1141-6545, it is usually assumed that $P_0 \ll P$, so that τ_{sd} reduces to the familiar characteristic age $\tau_c = P/(2\dot{P}) \simeq 1.5$ Myr. For the older recycled pulsars, however, Arzoumanian et al. (1999) pointed out that this assumption is usually not appropriate, since the weaker magnetic fields of these objects mean that their present spin periods are only moderately larger than the periods produced during accretion. Adopting the spin-up line from Arzoumanian et al. (1999), we may write

$$P_0 = \left(\frac{\dot{P}}{1.1 \times 10^{-15}} \right)^{3/4} \text{ s}. \quad (3)$$

Using the above two equations, we calculate τ_{sd} for the two recycled pulsars PSR J0751+1807 and PSR J1757-5322 to be 6.7 and 4.9 Gyr, respectively (see Table 1).

The remaining lifetime of a pulsar binary is defined by the shorter of the merger time of the binary due to the emission of GWs, τ_{mrg} , and the time that the pulsar will reach the “death line,” τ_d (Ruderman & Sutherland 1975). For young pulsars like PSR J1141-6545, which have relatively short radio lifetimes, $\tau_d < \tau_{\text{mrg}}$. Recycled pulsars, on the other hand, have far smaller spin-down rates than young pulsars, so it is likely that close binaries containing a recycled pulsar will coalesce before the pulsar reaches the death-line ($\tau_{\text{mrg}} < \tau_d$). For circular orbits, the results of the Peters (1964) calculations for the merger time of a binary system of two point masses m_1 and m_2 with orbital period P_b can be written as simply

$$\tau_{\text{mrg}} = 9.83 \times 10^6 \text{ yr} \left(\frac{P_b}{\text{hr}} \right)^{8/3} \left(\frac{\mu}{M_\odot} \right)^{-1} \left(\frac{m_1 + m_2}{M_\odot} \right)^{-2/3}, \quad (4)$$

where the reduced mass $\mu = m_1 m_2 / (m_1 + m_2)$. For the eccentric binary PSR J1141-6545, we use the more detailed calculations of Peters (1964) to calculate τ_{mrg} . Most of observed coalescing NS-WD binaries, as well as the DNS systems, have $\tau_{\text{mrg}} \sim 10^8 - 10^9$ yr.

Our understanding of pulsar emission is rather poor, and therefore, it is not clear how to calculate an accurate time associated with the termination of pulsar emission, and hence τ_d . Here we assume the spin-down torque is dominated by magnetic-dipole radiation with no evolution of the magnetic field. The surface magnetic field of a neutron star, B_s , can be estimated from the current spin period P (in s) and spin-down rate \dot{P} (in s s⁻¹):

$$B_s = 3.2 \times 10^{19} (P\dot{P})^{1/2} \text{ G}. \quad (5)$$

Chen & Ruderman (1993) comprehensively discussed the evolution of a pulsar period based on different magnetic field structures. Their results are consistent with previous studies (Ruderman & Sutherland 1975; van den Heuvel 1987). We adopt their case C (eq. [9] in their paper), according to which the radio emission terminates when the “death period”

$$P_d = \left(\frac{B_s}{1.4 \times 10^{11} \text{ G}} \right)^{7/13} \text{ s} \quad (6)$$

is reached. Assuming that the surface magnetic field remains constant, we can integrate equation (5) to calculate the time for the pulsar period to reach P_d . We find that

$$\tau_d = \left(\frac{P_d^2 - P^2}{2P\dot{P}} \right). \quad (7)$$

For PSR J1141–6545, we use equations (6) and (7) to find the remaining observable lifetime $\tau_d \sim 104$ Myr. This is significantly less than τ_{mrg} for this binary system (~ 600 Myr). Including the modest contribution from the characteristic age of PSR J1141–6545, we take the observable lifetime of the binary system to be $\tau_{\text{life}} = \tau_c + \tau_d \sim 105$ Myr. We note in passing that Edwards & Bailes (2001) estimated the remaining lifetime of PSR J1141–6545 to be only ~ 10 Myr. Although no details of their calculation were presented in their paper, they probably assumed some decay of the magnetic field, which led to their lower value of τ_d and hence τ_{life} .

In the cases of the recycled pulsars PSR J0751+1807 and PSR J1757–5322, which have lower magnetic field strengths and hence longer radio lifetimes, both binaries will coalesce before the pulsars stop radiating (i.e., $\tau_{\text{mrg}} \ll \tau_d$), so we calculate their lifetime using $\tau_{\text{life}} = \tau_{\text{sd}} + \tau_{\text{mrg}}$. The estimated lifetimes are 14.3 Gyr (PSR J0751+1807) and 12.7 Gyr (PSR J1757–5322), about 2 orders of magnitude longer than for the young NS-WD PSR J1141–6545.

2.2. Probability Density Function of the Galactic NS-WD Coalescence Rate Estimates

The basic strategy we use to calculate $P(\mathcal{R}_{\text{tot}})$ is described in detail in Paper I. In brief, using a detailed Monte Carlo simulation, for each observed NS-WD binary, we determine the fraction of the population that is actually *detectable* by careful modeling of all large-scale pulsar surveys. We include the selection effects that reduce the detectability of short-period binary systems when integration times are significant in comparison. Our pulsar population model takes into account the distribution of pulsars in the Galaxy and their luminosity function. Treating each pulsar separately, our simulations effectively probe the specific pulsar sub-populations with pulse and orbital characteristics similar to those of PSR J0751+1807, PSR J1757–5322, or PSR J1141–6545.

From the simulations we obtain N_{obs} pulsars detected by the surveys out of a Galactic population of N_{tot} in each model. We calculate N_{obs} repeatedly for a fixed N_{tot} . As shown in detail in Paper I, the distribution of N_{obs} follows a Poisson function, $P(N_{\text{obs}}; \langle N_{\text{obs}} \rangle)$. We calculate the best-fit value of $\langle N_{\text{obs}} \rangle$, which is the mean number of observed pulsars in a given sample, for a given N_{tot} . Here, N_{obs} is set to be unity, since each observed pulsar is unique and we consider them separately in our calculations. Thus, the likelihood of detecting *one pulsar* similar to the observed one from the given pulsar population with N_{tot} samples is simply $P(1; \langle N_{\text{obs}} \rangle)$. We vary N_{tot} and calculate $P(1; \langle N_{\text{obs}} \rangle)$ to determine the most probable value of N_{tot} . In addition, we found that $\langle N_{\text{obs}} \rangle$ is directly proportional to N_{tot} . We calculate α , which is the slope of the function $\langle N_{\text{obs}} \rangle = \alpha N_{\text{tot}}$ for each observed system for a pulsar population model.

Then, using Bayes's theorem, we calculate $P(\langle N_{\text{obs}} \rangle)$ from the likelihood $P(1; \langle N_{\text{obs}} \rangle)$ and eventually calculate $P(\mathcal{R})$ using a change of variables. We repeat the whole procedure for all three observed coalescing NS-WD binaries and combine the three individual PDFs to obtain a total PDF of Galactic coalescence rate of NS-WD binaries, $P(\mathcal{R}_{\text{tot}})$.

In Paper I, we showed that a normalized PDF of the coalescence rate for an individual pulsar binary system can be written as

$$P_i(\mathcal{R}) = C_i^2 \mathcal{R} e^{-C_i \mathcal{R}}, \quad (8)$$

where C_i is a coefficient determined by properties of the i th pulsar:

$$C_i \equiv \left(\frac{\alpha \tau_{\text{life}}}{f_b} \right)_i. \quad (9)$$

Here, the beaming correction factor f_b is the inverse of the fraction of 4π sr covered by the pulsar radiation beam during each rotation. In the case of the two DNS systems, PSR B1913+16 and PSR B1534+12, Kalogera et al. (2001) adopted $f_b \sim 6$ based on pulse profile and polarization measurements of two pulsars. The lack of such observations for the current sample of NS-WD binaries means that it is difficult to estimate reliable values of f_b . Therefore, in this paper, we do not correct for pulsar beaming (i.e., $f_b = 1$). As a result, all our values should be considered lower limits.

In Paper I, we calculated $P(\mathcal{R}_{\text{tot}})$ considering two observed DNS systems (labeled by the subscripts 1 and 2). We defined the total rate $\mathcal{R}_+ \equiv \mathcal{R}_1 + \mathcal{R}_2$ and showed that

$$P(\mathcal{R}_+) = \left(\frac{C_1 C_2}{C_2 - C_1} \right)^2 \left[\mathcal{R}_+ (e^{-C_1 \mathcal{R}_+} + e^{-C_2 \mathcal{R}_+}) - \left(\frac{2}{C_2 - C_1} \right) (e^{-C_1 \mathcal{R}_+} - e^{-C_2 \mathcal{R}_+}) \right], \quad (10)$$

where $C_1 < C_2$. In Appendix A, we show that this can be extended for the current case of interest, in which we have three binary systems such that $\mathcal{R}_+ \equiv \mathcal{R}_1 + \mathcal{R}_2 + \mathcal{R}_3$. This leads to

$$\begin{aligned} P(\mathcal{R}_+) = & \frac{C_1^2 C_2^2 C_3^2}{(C_2 - C_1)^3 (C_3 - C_1)^3 (C_3 - C_2)^3} \\ & \times \left((C_3 - C_2)^3 e^{-C_1 \mathcal{R}_+} \{ -2(-2C_1 + C_2 + C_3) + \mathcal{R}_+ [-C_1(C_2 + C_3) + (C_1^2 + C_2 C_3)] \} \right. \\ & + (C_3 - C_1)^3 e^{-C_2 \mathcal{R}_+} \{ 2(C_1 - 2C_2 + C_3) + \mathcal{R}_+ [C_2(C_3 + C_1) - (C_2^2 + C_3 C_1)] \} \\ & \left. + (C_2 - C_1)^3 e^{-C_3 \mathcal{R}_+} \{ -2(C_1 + C_2 - 2C_3) + \mathcal{R}_+ [-C_3(C_1 + C_2) + (C_3^2 + C_1 C_2)] \} \right), \quad (11) \end{aligned}$$




FIG. 1.—PDFs of the Galactic coalescence rate estimation in both a logarithmic and a linear scale (*inset*) are shown for the reference model. The solid line represents $P(\mathcal{R}_{\text{tot}})$. Other curves are $P(\mathcal{R})$ for PSR J1757–5322 (*dot-dashed line*), PSR J0751+1807 (*short-dashed line*), and PSR J1141–6545–like populations (*long-and-short dashed lines*), respectively. Dotted lines correspond to 68%, 95%, and 99% confidence limits for $P(\mathcal{R}_{\text{tot}})$.

where the coefficients C_i ($i = 1, 2, 3$) are defined by equation (9) and $C_1 < C_2 < C_3$. This result was already used in our recent rate estimation for DNS systems to include the newly discovered pulsar, PSR J0737–3039 (Burgay et al. 2003; Kalogera et al. 2004). As before, the confidence limit (CL)⁴ and the lower and upper limits (\mathcal{R}_l and \mathcal{R}_u) of the coalescence rate estimates are defined in the same way we described in Paper I; i.e.,

$$\int_{\mathcal{R}_l}^{\mathcal{R}_u} P(\mathcal{R}_+) d\mathcal{R}_+ = \text{CL}, \quad (12)$$

and

$$P(\mathcal{R}_l) = P(\mathcal{R}_u). \quad (13)$$

3. RESULTS

In Figure 1, we show the resulting $P(\mathcal{R}_{\text{tot}})$ for NS-WD binaries along with the individual PDFs for each observed coalescing binaries. The figure shown here is obtained from our reference model (model 6 in Paper I). As we found for the DNS systems in Paper I, $P(\mathcal{R}_{\text{tot}})$ is highly peaked and dominated by a single object. In this case, PSR J1141–6545 dominates the results by virtue of its short observable lifetime ($\tau_{\text{life}} \sim 105$ Myr). This is in spite of the fact that the estimated total number of binaries similar to PSR J0751+1807 ($N_{0751} \simeq 2900$) is the largest among the observed systems.

⁴ Strictly speaking, the phrase “confidence limit” used in this paper should be interpreted as a confidence *interval* such that a true value of the Galactic coalescence rate would exist in a given range of rates. However, we keep the terminology “confidence limit” for consistency with Paper I and Kalogera et al. (2004).

TABLE 2
ESTIMATES FOR THE GALACTIC COALESCENCE RATE (\mathcal{R}_{tot}) OF NS-WD BINARIES
AT VARIOUS CONFIDENCE LIMITS FOR ALL MODELS CONSIDERED

MODEL ^a	\mathcal{R}_{tot} (Myr ⁻¹)		
	Peak ^b	68% ^c	95% ^c
1.....	1.23	+1.57 -0.77	+3.97 -1.04
2.....	1.00	+1.27 -0.62	+3.20 -0.83
3.....	1.32	+1.69 -0.83	+4.27 -1.12
4.....	1.53	+1.99 -0.97	+5.02 -1.31
5.....	1.19	+1.53 -0.75	+3.87 -1.01
6.....	4.11	+3.23 -2.56	+13.23 -3.47
7.....	1.73	+2.21 -1.08	+5.57 -1.46
8.....	0.83	+1.07 -0.52	+2.71 -0.71
9.....	0.43	+0.55 -0.27	+1.39 -0.36
10.....	1.42	+1.88 -0.92	+4.76 -1.23
11.....	0.72	+0.96 -0.47	+2.42 -0.62
12.....	0.55	+0.72 -0.35	+1.82 -0.47
13.....	0.40	+0.53 -0.26	+1.33 -0.34
14.....	0.23	+0.30 -0.15	+0.76 -0.20
15.....	10.03	+12.46 -6.09	+31.38 -8.34
16.....	3.68	+4.57 -2.25	+11.53 -3.07
17.....	2.45	+3.06 -1.50	+7.71 -2.04
18.....	1.55	+1.94 -0.95	+4.90 -1.29
19.....	0.72	+0.91 -0.44	+2.29 -0.60
20.....	5.48	+6.69 -3.29	+16.86 -4.52
21.....	1.96	+2.37 -1.17	+5.96 -1.61
22.....	1.55	+1.97 -0.96	+4.96 -1.30
23.....	1.13	+1.47 -0.71	+3.72 -0.97
24.....	1.14	+1.49 -0.73	+3.76 -0.98
25.....	1.21	+1.56 -0.76	+3.95 -1.03
26.....	1.33	+1.72 -0.84	+4.35 -1.13
27.....	1.46	+1.90 -0.93	+4.79 -1.25

^a Model number (see Paper I for details of model parameters).

^b Peak value of the probability density function.

^c Confidence limits.

We summarize our results for different pulsar population models in Table 2. The model parameters are identical to those described in Paper I. We note, however, following Kalogera et al. (2004), that our reference model is now model 6 ($L_{\text{min}} = 0.3$ mJy kpc²) rather than model 1 ($L_{\text{min}} = 1.0$ mJy kpc²). This choice reflects the recent discoveries of faint pulsar with 1400 MHz radio luminosities below 1.0 mJy kpc² (Camilo 2003). The peak values of $P(\mathcal{R}_{\text{tot}})$ lie in the range between ~ 0.2 and 10 Myr⁻¹, where the reference model shows a peak around 4 Myr⁻¹. For the reference model, the uncertainties in the rates (defined by $\mathcal{R}_u/\mathcal{R}_l$) are estimated to be ~ 6 , 27 , and 62 at 68%, 95%, and 99% CL, respectively. Comparing this to results from Kalogera et al. (2004), we find that the uncertainties at different CL of the coalescence rate of NS-WD binaries are typically larger by factor of ~ 1.4 than those of the DNS systems. This result is robust for all models we consider.

The correlations between the peak value of the total Galactic coalescence rate $\mathcal{R}_{\text{peak}}$ and the model parameters (e.g., the cutoff luminosity L_{min} and the power index p) seem to be similar to those of DNSs we observed in Paper I. As we found for the DNS systems, $\mathcal{R}_{\text{peak}}$ values strongly depend on a pulsar luminosity function rather than a spatial distribution of pulsars in the Galaxy; in other words, $\mathcal{R}_{\text{peak}}$ values rapidly increase as the fraction of faint pulsars increases.

4. GRAVITATIONAL WAVE BACKGROUND DUE TO NS-WD BINARIES

Close binaries consisting of compact objects (e.g., NS-WD binaries) are suggested as important GW sources in a frequency range below 1 mHz. In this range, because of the large number of sources, *LISA* would not be able to resolve each source within a given frequency band. Hence the Galactic binaries are expected to establish a confusion noise level (or “background”) dominated by WD-WD binaries (Hils et al. 1990; Bender & Hils 1997; Nelemans et al. 2001; Schneider et al. 2001). In this work, we consider the contribution from NS-WD binaries to the predicted confusion noise level. Using our results from the previous section, we calculate the amplitude of GW signals from NS-WD binaries in the nearby universe and compare it with the *LISA* sensitivity curve.⁵ We assume that the three observed systems represent the whole population of NS-WD binaries in our Galaxy.

We calculate the characteristic strain amplitude of GWs (h_c) from NS-WD binaries using the results given by Phinney (2001) for circular binaries. In general, binaries with an eccentricity e emit GWs at frequencies $f = n\nu$, i.e., the n th harmonic of the orbital frequency ν . In the case of circular binaries, $n = 2$ because of the orbital symmetry and the quadrupole nature of GWs. The

⁵ We use the online sensitivity curve generator to calculate the sensitivity curve of *LISA* (<http://www.srl.caltech.edu/~shane/sensitivity/MakeCurve.html>).

eccentricity of PSR J0751+1807 and PSR J1757–5322 are $e \sim 10^{-4}$ and $\sim 10^{-6}$, respectively. Hence, it is safe to consider them as circular binaries. PSR J1141–6545 has an appreciable eccentricity ($e = 0.17$), but for simplicity, we consider only the $n = 2$ harmonic as if it were a circular binary.⁶

For an observation of length T_{obs} with a GW detector, the contribution from background sources (NS-WD binaries in this work) depends on the number of sources within the frequency resolution, $\Delta f = 1/T_{\text{obs}}$. Following Schneider et al. (2001), we define an effective GW amplitude $h_{\text{rms}}(f) \equiv h_c(f)(\Delta f/f)^{1/2}$, where $h_c(f)$ is a characteristic strain amplitude. Phinney (2001) showed a simple analytic formula to calculate $h_c(f)$ for a population of inspiraling circular-orbit binaries with a given number density in the nearby universe. We use equation (16) in his paper to calculate $h_c(f)$,⁷ and we find

$$h_{\text{rms}}(f) \simeq 1.7 \times 10^{-26} \left(\frac{\mathcal{M}}{M_{\odot}} \right)^{5/6} \left(\frac{f}{\text{mHz}} \right)^{-7/6} \left(\frac{N_0}{\text{Mpc}^{-3}} \right)^{1/2} \left(\frac{T_{\text{obs}}}{\text{yr}} \right)^{-1/2}, \quad (14)$$

where \mathcal{M} is the ‘‘chirp mass’’ of an NS-WD binary, defined by

$$\mathcal{M} \equiv \frac{(M_{\text{NS}} M_{\text{WD}})^{3/5}}{(M_{\text{NS}} + M_{\text{WD}})^{1/5}}, \quad (15)$$

and N_0 is the comoving number density of NS-WD, i.e., the number of sources per Mpc^3 .

We calculate the GW amplitude of NS-WD binaries in a frequency range $f_{\text{min}} < f < f_{\text{max}}$. Estimated GW frequencies of three NS-WD binaries based on their current separations are all less than ~ 0.1 mHz. In our calculation, however, we set the minimum frequency f_{min} to be 1 mHz taking into account the fact that the confusion noise level is mainly dominated by Galactic WD-WD binaries at lower frequency range $f < 1$ mHz (Nelemans et al. 2001). The maximum GW frequency f_{max} is calculated by $f_{\text{max}} = 2/P_{b,\text{min}}$, where $P_{b,\text{min}}$ is the minimum orbital period of the binary at the WD Roche-lobe overflow.

Following Eggleton (1983), we calculate the minimum possible separation of the binary using $a_{\text{min}} = R_{\text{WD}}/r_L$, where r_L is the effective Roche lobe radius. We estimate the radius of a white dwarf companion R_{WD} adopting the results given by Tout et al. (1997). We show the estimated mass of white dwarf companions in Table 1.

Converting a_{min} to f_{max} based on Kepler’s third law, we find that

$$f_{\text{max}} \simeq 0.16 r_L^{3/2} \left(\frac{M_{\text{WD}} + M_{\text{NS}}}{M_{\odot}} \right)^{1/2} \left[\left(\frac{M_{\text{ch}}}{M_{\text{WD}}} \right)^{2/3} - \left(\frac{M_{\text{WD}}}{M_{\text{ch}}} \right)^{2/3} \right]^{-3/4} \text{ Hz}, \quad (16)$$

where $M_{\text{ch}} = 1.44 M_{\odot}$ is the Chandrasekhar limit. We note that $f_{\text{max}} = f_{\text{max}}(M_{\text{NS}}, M_{\text{WD}})$. Based on equation (16), we define three frequency regions: (a) $f_{\text{min}} < f < f_{\text{max},0751}$, (b) $f_{\text{max},0751} < f < f_{\text{max},1757}$, and (c) $f_{\text{max},1757} < f < f_{\text{max},1141}$. In region *a*, for example, all three observed NS-WD systems contribute to the GW background. However, for frequencies larger than $f_{\text{max},0751}$, PSR J0751+1807–like populations have already reached the Roche lobe overflow and we can not apply equation (14) to these systems. Therefore, in a frequency region *b*, we consider PSR J1141–6545–like and PSR J1757–5322–like populations. Similarly, for the highest frequency range (region *c*), we consider the contribution to the GW signals from PSR J1141–6545–like population only.

In order to calculate $h_{\text{rms}}(f)$, we need the chirp mass \mathcal{M} and present-day comoving number density of NS-WD binaries N_0 . Following Farmer & Phinney (2003), we define the ‘‘flux-weighted’’ averaged chirp mass:

$$\langle \mathcal{M} \rangle \equiv \frac{\sum F_{\text{gw},i} \mathcal{M}_i}{\sum F_{\text{gw},i}} = \frac{\sum N_{\text{peak},i} \mathcal{M}_i^{13/3}}{\sum N_{\text{peak},i} \mathcal{M}_i^{10/3}}, \quad (17)$$

where \mathcal{F}_{gw} is the GW flux ($\mathcal{F}_{\text{gw}} \propto f^{10/3} \mathcal{M}^{10/3} \mathcal{N}_{\text{peak}}$) and $\mathcal{N}_{\text{peak}}$ is the peak value of $P(N_{\text{tot}})$ of the each subpopulation of NS-WD binaries in our Galaxy. The subscript *i* represents each pulsar subpopulation.

For example, in region *b*, $\langle \mathcal{M} \rangle = (\mathcal{N}_{1757} \mathcal{M}_{1757}^{13/3} + \mathcal{N}_{1141} \mathcal{M}_{1141}^{13/3}) / (\mathcal{N}_{1757} \mathcal{M}_{1757}^{10/3} + \mathcal{N}_{1141} \mathcal{M}_{1141}^{10/3})$. Because $\mathcal{N}_{\text{peak}}$ is a constant and independent of the GW frequency, it follows that $\langle \mathcal{M} \rangle$ is independent of frequency. As a result, the evolution of orbital characteristics and hence the GW frequency of a binary are solely determined by the inspiral process. This is true regardless of the initial distribution of orbital characteristics.

We now calculate the comoving number density of NS-WD binaries

$$N_0 = \int_0^{\infty} N(z) dz, \quad (18)$$

⁶ We calculate the power distribution in various harmonics for this eccentricity based on the result of Peters & Mathews (1963). We note that the GW amplitude calculated for PSR J1141–6545 in this work corresponds to $\sim 70\%$ of the total power of the gravitational radiation emitted from this binary. The remaining power is contributed from the higher harmonics.

⁷ We assume a case in which the last term in Phinney’s eq. (16), $[(1+z)^{-1/3}/0.74]^{1/2}$, becomes unity. The calculation is not significantly affected by different assumptions of cosmological models and comoving number density functions of coalescing binaries (Phinney 2001).




FIG. 2.—Effective GW amplitude h_{rms} for coalescing NS-WD binaries overlapped with the *LISA* sensitivity curve. The curve is produced with the assumption of $S/N = 1$ for 1 yr of integration. Dotted lines are results from all models we consider except the reference model, which is shown as a solid line. We also show the expected confusion noise from Galactic WD-WD binaries for comparison (*dashed line*).

where $N(z) dz$ is the number of NS-WD binaries per unit comoving volume between redshift z and $z + dz$. Noting that the number density of NS-WD binaries is proportional to the total number of systems, we may write

$$N_0 = \epsilon \mathcal{N}_{\text{PSR}}, \quad (19)$$

where ϵ is the star formation rate density per unit comoving volume ($\dot{\rho}$) normalized to the Galactic star formation rate (r); i.e., $\epsilon = \dot{\rho}/r$ (in Mpc^{-3}). \mathcal{N}_{PSR} is the most likely value of the total number of pulsars for each frequency range (e.g., $\mathcal{N}_{\text{PSR}} = \mathcal{N}_{0751} + \mathcal{N}_{1757} + \mathcal{N}_{1141}$ in region *a*). We derived $P(N_{\text{tot}})$ for individual systems in Paper I. In a similar fashion to the coalescence rate estimation described in Appendix A, the combined PDF of N_{tot} can be calculated from individual PDFs of the observed systems. Then \mathcal{N}_{PSR} can be obtained from the peak value of the PDF we calculate for each frequency range we discussed above.

Using the results of Cole et al. (2001), we find

$$\epsilon = \frac{\dot{\rho}(0)}{r} \int_0^{z_{\text{max}}} \frac{[1 + (b/a)z]}{[1 + (z/c)^d]} dz \text{ Mpc}^{-3}, \quad (20)$$

where $(a, b, c, d) = (0.0166, 0.1848, 1.9474, 2.6316)$ are parameters that take into account dust-extinction corrections (see Cole et al. (2001) for further details). Assuming a Hubble constant $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$, we calculate the Galactic star formation rate density $\dot{\rho}(0) \simeq 0.01 M_{\odot} \text{ yr}^{-1} \text{ Mpc}^{-3}$. Following Cappellaro et al. (1999), assuming the Salpeter initial mass function, we convert the Galactic supernova Type II+Ib/c rate⁸ to the star formation rate and find $r \sim 0.7 M_{\odot} \text{ yr}^{-1}$. Numerically integrating

⁸ Cappellaro et al. (1999) assumed a Hubble constant $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$. Since we adopt $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$, we have multiplied their results by a factor of $(65/75)^2$. We also note that we consider Sbc–Sd type galaxies only, which would be relevant for active star-forming regions.

equation (20) out to $z_{\max} = 5$, which is considered to be the onset of the galaxy formation (Schneider et al. 2001), we find $\epsilon \sim 0.6 \text{ Mpc}^{-3}$. The number density of NS-WD binaries N_0 then can be calculated by equation (19) for a given \mathcal{N}_{PSR} for each model.

In Figure 2, we plot the GW amplitude h_{rms} against the simulated *LISA* sensitivity curve calculated for a signal-to-noise ratio $S/N = 1$. All dotted lines correspond to the GW amplitude calculated from the full set of pulsar population models we consider and the solid line is the result from our reference model. The range of the GW amplitude for all models spans about an order of magnitude. We find that the GW background amplitude from NS-WD binaries is about 1–2 orders of magnitude smaller than the expected sensitivity curve of *LISA* at GW frequencies larger than 1 mHz and it is unlikely that this population will be detected with *LISA*. In the lower frequency region (below ~ 1 mHz), the GW background amplitude from NS-WD binaries increases as f decreases. However, the contribution from NS-WD binaries to the GW background noise level would still be less than $\sim 10\%$ of the GW amplitude from WD-WD binaries (Fig. 2, *dashed line*). We note that, however, we have not considered any beaming corrections, so the NS-WD background curves should be viewed as lower limits. This possibility is discussed briefly in the next section.

5. DISCUSSION

We have used detailed Monte Carlo simulations to calculate the Galactic coalescence rate of NS-WD binaries. From the reference model, the most probable value of \mathcal{R}_{tot} is estimated to be $4.11_{-2.56}^{+5.25} \text{ Myr}^{-1}$ at a 68% statistical confidence limit. We find that the coalescence rate of NS-WD binaries is about factor of 20 smaller than those of DNS for all pulsar population models we consider. As mentioned above, we did not take into account any beaming correction for NS-WD binaries. If we assume a beaming fraction of pulsars in NS-WD binaries similar to that of pulsars found in DNS, $f_b \sim 6$, then the discrepancy between $\mathcal{R}_{\text{peak,DNS}}$ and $\mathcal{R}_{\text{peak,NS-WD}}$ is significantly reduced. As a simple estimate, if we assume $f_{b,1141} \sim 5$, but keeping $f_b = 1$ for the other two binaries, the estimated Galactic coalescence rate increases to $18.06_{-12.74}^{+26.05} \text{ Myr}^{-1}$ at a 68% confidence limit. Hence the ratio between DNS and NS-WD coalescence rate decreases to about 5. Because the contribution from PSR J0751+1807 and PSR J1757–5322 is an order of magnitude smaller than that of PSR J1141–6545, moderate values of beaming fraction for those recycled pulsars do not change the result significantly.

Based on the number of sources of NS-WD binaries in our Galaxy, we estimate the effective GW amplitude from the cosmic population of these systems. We find that the GW background from NS-WD binaries is too weak to be detected by *LISA* for the nominal beaming correction. Only by adopting an unreasonably large beaming correction factor, $f_b > 10$, could these systems be detectable by *LISA* in the mHz range. These results are in good agreement with an independent study by Cooray (2004) based on statistics of low-mass X-ray binaries.

We finally note that combining the results from Paper I and this work can give us strong constraints on the population synthesis models. The preferred models, which are consistent with both $\mathcal{R}_{\text{NS-WD}}$ and \mathcal{R}_{DNS} , can then be used for the estimation of the coalescence rate of neutron star–black hole binaries, which have not yet been observed.

We thank A. Cooray for noticing the frequency resolution correction factor in the calculation of the GW background. V. K. would like to acknowledge partial support from NSF grant PHY 01-21420 and a Packard Fellowship in Science and Engineering to V. K., and D. R. L. is a University Research Fellow funded by the Royal Society. He is also grateful for support from the Theoretical Astrophysics Visitors' Fund at Northwestern University.

APPENDIX A

COMBINED RATE PDF FOR THREE BINARY SYSTEMS

In Paper I, we derived expressions for $P(\mathcal{R}_{\text{tot}})$ for one and two coalescing binaries. In this paper, and in our revised DNS coalescence rate estimates (Kalogera et al. 2004), we extend this PDF to the case of three systems. Following Paper I, we define a coefficient for each observed NS-WD:

$$A \equiv \left(\frac{\alpha \tau_{\text{life}}}{f_b} \right)_{1141}, \quad B \equiv \left(\frac{\alpha \tau_{\text{life}}}{f_b} \right)_{0751}, \quad \text{and} \quad C \equiv \left(\frac{\alpha \tau_{\text{life}}}{f_b} \right)_{1757}, \quad (\text{A1})$$

where $A < B < C$. Recall that α is the slope of the function $\langle N_{\text{obs}} \rangle = \alpha N_{\text{tot}}$ and is determined for each pulsar population model for each NS-WD system. By definition, the total Galactic coalescence rate is the sum of all three observed systems:

$$\mathcal{R}_+ \equiv \mathcal{R}_1 + \mathcal{R}_2 + \mathcal{R}_3. \quad (\text{A2})$$

Redefining $\mathcal{R}_+ \equiv \mathcal{R}_a + \mathcal{R}_b$, where $\mathcal{R}_a \equiv \mathcal{R}_1 + \mathcal{R}_2$ and $\mathcal{R}_b \equiv \mathcal{R}_3$, we transform \mathcal{R}_a and \mathcal{R}_b to new variables \mathcal{R}_+ and $\mathcal{R}_- \equiv \mathcal{R}_a - \mathcal{R}_b$:

$$P(\mathcal{R}_+, \mathcal{R}_-) = P(\mathcal{R}_a, \mathcal{R}_b) \begin{vmatrix} \frac{d\mathcal{R}_a}{d\mathcal{R}_+} & \frac{d\mathcal{R}_b}{d\mathcal{R}_-} \\ \frac{d\mathcal{R}_a}{d\mathcal{R}_-} & \frac{d\mathcal{R}_b}{d\mathcal{R}_+} \end{vmatrix} = \frac{1}{2} P(\mathcal{R}_a, \mathcal{R}_b). \quad (\text{A3})$$

Since both \mathcal{R}_+ and \mathcal{R}_- are positive, $-\mathcal{R}_- \leq \mathcal{R}_+ \leq +\mathcal{R}_-$.

The PDF of the total rate \mathcal{R}_+ is obtained after integrating $P(\mathcal{R}_+, \mathcal{R}_-)$ over \mathcal{R}_- :

$$P(\mathcal{R}_+) = \int_{\mathcal{R}_-} P(\mathcal{R}_+, \mathcal{R}_-) d\mathcal{R}_- = \frac{1}{2} \int_{\mathcal{R}_-} P(\mathcal{R}_a, \mathcal{R}_b) d\mathcal{R}_-, \quad (\text{A4})$$

where

$$P(\mathcal{R}_a, \mathcal{R}_b) = P(\mathcal{R}_a)P(\mathcal{R}_b). \quad (\text{A5})$$

Here, $P(\mathcal{R}_a = \mathcal{R}_1 + \mathcal{R}_2)$ is given by equation (10) and we can rewrite the formula with appropriate coefficients defined earlier:

$$P(\mathcal{R}_a) = \left(\frac{AB}{B-A} \right)^2 \left[\mathcal{R}_a (e^{-A\mathcal{R}_a} + e^{-B\mathcal{R}_a}) - \left(\frac{2}{B-A} \right) (e^{-A\mathcal{R}_a} - e^{-B\mathcal{R}_a}) \right]. \quad (\text{A6})$$

The individual PDF $P(\mathcal{R}_b) \equiv P(\mathcal{R}_3)$ (eq. [8]) is also rewritten as follows:

$$P_i(\mathcal{R}_b) = C^2 \mathcal{R}_b e^{-C\mathcal{R}_b}. \quad (\text{A7})$$

Replacing $\mathcal{R}_a = (1/2)(\mathcal{R}_+ + \mathcal{R}_-)$ and $\mathcal{R}_b = (1/2)(\mathcal{R}_+ - \mathcal{R}_-)$ in equation (A4), the normalized $P(\mathcal{R}_+)$ can be obtained by integration. After some algebra, we find

$$P(\mathcal{R}_+) = \frac{A^2 B^2 C^2}{(B-A)^3 (C-A)^3 (C-B)^3} \times \left((C-B)^3 e^{-A\mathcal{R}_+} \{ -2(-2A+B+C) + \mathcal{R}_+ [-A(B+C) + (A^2+BC)] \} \right. \\ \left. + (C-A)^3 e^{-B\mathcal{R}_+} \{ 2(A-2B+C) + \mathcal{R}_+ [B(C+A) - (B^2+CA)] \} \right. \\ \left. + (B-A)^3 e^{-C\mathcal{R}_+} \{ -2(A+B-2C) + \mathcal{R}_+ [-C(A+B) + (C^2+AB)] \} \right). \quad (\text{A8})$$

REFERENCES

- Abramovici, A., et al. 1992, *Science*, 256, 325
 Arzoumanian, Z., Cordes, J. M., & Wasserman, I. 1999, *ApJ*, 520, 696
 Bailes, M., Ord, S. M., Knight, H. S., & Hotan, A. W. 2003, *ApJ*, 595, L49
 Bender, P. L., & Hils, D. 1997, *Classical Quantum Gravity*, 14, 1439
 Bender, P., et al. 1998, *LISA Pre-Phase A Report* (2nd ed.; Garching: Max-Planck-Institut für Quantenoptik)
 Burgay, M., et al. 2003, *Nature*, 426, 531
 Camilo, F. 2003, in *ASP Conf. Ser. 302, Radio Pulsars*, ed. M. Bailes, D. J. Nice, & S. E. Thorsett (San Francisco: ASP), 145
 Cappellaro, E., Evans, R., & Turatto, M. 1999, *A&A*, 351, 459
 Chen, K., & Ruderman, M. 1993, *ApJ*, 402, 264
 Cole, S., et al. 2001, *MNRAS*, 326, 255
 Cooray, A. 2004, *MNRAS*, 354, 25
 Curran, S. J. & Lorimer, D. R. 1995, *MNRAS*, 276, 347
 Edwards, R. T., & Bailes, M. 2001, *ApJ*, 547, L37
 Eggleton, P. P. 1983, *ApJ*, 268, 368
 Farmer, A. J., & Phinney, E. S. 2003, *MNRAS*, 346, 1197
 Hils, D., Bender, P. L., & Webbink, R. F. 1990, *ApJ*, 360, 75
 Kalogera, V., Narayan, R., Spiegel, D. N., & Taylor J. H. 2001, *ApJ*, 556, 340
 Kalogera, V., et al. 2004, *ApJ*, 601, L179 (erratum 614, L137)
 Kaspi, V. M., et al. 2000, *ApJ*, 543, 321
 Kim, C., Kalogera, V., & Lorimer, D. 2003, *ApJ*, 584, 985 (Paper I)
 Larson, S. L., Hiscock W. A., & Hellings, R. W. 2000, *Phys. Rev. D*, 62, 062001
 Lorimer, D. R. 2001, *Binary and Millisecond Pulsars at the New Millennium* (Potsdam: Max-Planck-Gesellschaft), <http://relativity.livingreviews.org/Articles/lrr-2001-5r>
 Lundgren, S. C., Zepka, A. F., & Cordes, J. M. 1995, *ApJ*, 453, 419
 Manchester, R. N., & Taylor, J. H. 1977, *Pulsars* (San Francisco: Freeman)
 Narayan, R. 1987, *ApJ*, 319, 162
 Narayan, R., Piran, T., & Shemi, A. 1991, *ApJ*, 379, L17
 Nelemans, G., Yungelson, L. R., & Portegies Zwart, S. F. 2001, *A&A*, 375, 890
 Nice, D. J., Splaver, E. M., & Stairs, I. H. 2004, *IAU*, 218, 49
 Peters, P. C. 1964, *Phys. Rev. B*, 136, 1224
 Peters, P. C., & Mathews, J. 1963, *Phys. Rev. D*, 131, 435
 Phinney, E. S. 1991, *ApJ*, 380, L17
 ———. 2001, preprint (astro-ph/0108028)
 Ruderman, M. A., & Sutherland, P. G. 1975, *ApJ*, 196, 51
 Schneider, R., Ferrari, V., Matarrese, S., & Portegies Zwart, S. F. 2001, *MNRAS*, 324, 797
 Tout, C. A., Aarseth, S. J., & Pols, O. R. 1997, *MNRAS*, 291, 732
 van den Heuvel, E. P. J. 1987, in *IAU Symp. 125, The Origin and Evolution of Neutron Stars*, ed. D. J. Helfand & J.-H. Huang (Dordrecht: Kluwer), 393