


Lectures 16-19

Properties of Ecosystems

- The flow of energy powers ecosystems.
- Matter cycles between the biotic and abiotic world.

Understanding Ecosystem Terms



Mean Residence Time

For a system in dynamic equilibrium:

Mean Residence Time (τ) = stock / inflow or outflow



τ = Average length of time a given atom or molecule spends in the system between entering and leaving.

Lectures 16-19


Importance of Nutrient Cycles

Hydrologic Cycle (water)

Read pages 36 – 37 in textbook

Hydrologic Cycle (water)

Pools (units are 10^6 km^3)



0.013


9.5

0.1


9.6

Lectures 16-19

Change in flow of Colorado River




Humans Alter the Hydrologic Cycle



From 1973 to 1987 the Aral dropped from fourth to sixth among the world's largest lakes.

Changes in Ogallala Aquifer 1980-1995




Lectures 16-19

Carbon Cycle


Reciprocal Processes:

- Photosynthesis (P_s)-
- Cellular Respiration (R) -

Active Carbon Pools




Global Carbon Cycle




Lectures 16-19

Atmospheric CO₂ was in dynamic equilibrium for thousands of years.




Change in atmospheric CO₂ concentration



The Breathing Earth

Jan. - Feb.




July - Aug.




Lectures 16-19

More land & pollution in N. Hemisphere



Seasonal Imbalances between Photosynthesis & Aerobic Respiration



Carbon Cycle Mean Residence Times

- Oceans ~ 422 years
- Atmosphere ~ 3 years (but atmospheric lifetime > 100 years)
- Land Plants ~ 4.6 years
- At current rates of fossil fuel use, our recoverable supplies will last about 700 years ! (4000 Pg ÷ 6.4 Pg/yr = 625 yrs)

Lectures 16-19

The Carbon Cycle is not Always in Balance

- In the present, inputs of CO_2 > outputs to the atmosphere because ...
 - Burning of fossil fuels.
 - Net destruction of terrestrial vegetation.
- Current imbalance results in rising atmospheric CO_2 concentrations.
- In the past, $P_s > R$ because ...
 - Fossil fuel deposits formed.
 - Oxygen accumulated in the atmosphere.

The Missing Carbon Sink

Net CO_2 Sources


Fossil Fuel Emission	6.4
Land Use Change	1.5
<hr/>	
Total Net Sources	7.9

Net CO_2 Sinks

Atmospheric Increase	3.2
Ocean uptake	2.2
<hr/>	
Total Net Sinks	5.4


Units are Pg C/yr

Mother nature mitigates
climate change to some extent.




Lectures 16-19

Historical estimates of sources & sinks of CO₂ associated with human activity



Unless we change, we are on the fast track



Summary of Carbon Cycle

- Largest active pool is the ocean.
- Pool on land (plant + soil) \cong 3x amount in atmosphere
- $P_s \cong R_{total}$
- Soil respiration \cong 10x fossil fuel emissions
- P_s removes 1/6 of the atmospheric pool of CO₂ each year.
- Concentration of CO₂ in the atmosphere is ca. 0.4% per year (ca. 3×10^{15} g C/yr)

Lectures 16-19

The Nitrogen (N) Cycle

- Atmosphere is ca. 78 % N_2 but most is unavailable to living things because ...
- N is important because ...
- Microbial processes are important in the steps of the N cycle.

The nitrogen cycle has 5 basic steps

- 1) Nitrogen Fixation: $N_2 \Rightarrow NH_3$

Root Nodules on a Legume



Lectures 16-19

Heterocysts in *Anabaena*



Heterocyst

Humans Fix Nitrogen Too !



Fritz Haber



at 500°C & several hundred atmospheres of pressure



2) Ammonification: organic N \Rightarrow NH_3

3) Nitrification: $\text{NH}_3 \Rightarrow \text{NO}_2^- \Rightarrow \text{NO}_3^-$

2-step process - each step by different bacteria.

Step 1: oxidation of ammonia (NH_3) to nitrite (NO_2^-) by _____


Step 2: oxidation of nitrite (NO_2^-) to nitrate (NO_3^-) by _____

Both steps couple E-releasing oxidations to fixation of carbon - chemoautotrophs.

Lectures 16-19

4) Nitrogen Assimilation: $\text{NH}_3 \Rightarrow$ organic N
 $\text{NO}_3^- \Rightarrow$ organic N


5) Denitrification: NO_3^- or $\text{NO}_2^- \Rightarrow \text{N}_2$ or N_2O




Active N Pools

- Atmosphere $3,800,000 \times 10^{15} \text{ g N}$
- Ocean $21,000 \times 10^{15} \text{ g N}$
- Soil Organic Matter $95 \times 10^{15} \text{ g N}$
- Terrestrial Biota $3.5 \times 10^{15} \text{ g N}$

Lectures 16-19



Human Activities Account for More than Half of Total N Inputs on Land



Terrestrial Ecosystems Can be Overfertilized

Potential Consequences of N Saturation

- Increased surface-water NO_3^- concentrations.
- Enhanced losses of nutrient cations.
- Soil acidification & greater soluble Al .


Summary of N Cycle


- Largest active pool = N_2 in atmosphere which is 181x > amount in ocean
- N in soil organic matter is 27x > amount in terrestrial biota
- Largest flux = uptake by plants of which almost all is from recycled organic N
- Human activities \approx 60 % of total inputs to land
- River flow \approx 20% total inputs to oceans


The Phosphorus (P) Cycle

- Example of a sedimentary cycle => no gaseous phase
- P is abundant in soil but in forms that are not readily available to biota
- PO_4^{3-} is an available form of P
- P is important because ...

Lectures 16-19








Lectures 16-19


Guano Islands



Mining guano ca. 1860.



The Global P Cycle



Lectures 16-19


Summary of P Cycle

- Abundant but low availability.
- Weathering of P-rich rock is original source.
- Geologic processes are slow (millions of years) so biota rapidly recycle organic-P.
- Residence time in biota is only a few days in the ocean.

Summary of P Cycle


- Large loss to ocean relative to rate of return to land.
- Losses in runoff are 90% particulate-P
- Mycorrhizae absorption by plant roots
- Mining P-rich rocks is a major source to land.

Aquatic Ecosystems Can be Overfertilized




Lectures 16-19

Temperate Lakes Can Thermally Stratify In The Summer



Eutrophication of stratified shallow lakes can change species composition



Cleaner water has allowed mayfly populations to increase again!



Summer 2012
Lakeside, Ohio



Lectures 16-19

Emerging Trends May Cause the return of Eutrophication to Lake Erie

Record-setting algal bloom in Lake Erie caused by agricultural and meteorological trends consistent with expected future conditions





Fig. 5. MODIS satellite image of Lake Erie on September 3, 2011, overlaid over map of Lake Erie tributaries. This image shows the bloom about 8 wk after its initiation in the western basin. On this date, it covers the entire western basin and is beginning to expand into the central basin, where it will continue to grow until October (Fig. 5f).

5448-5452 | PNAS | April 16, 2013 | vol. 110 | no. 16


Eutrophication in Coastal Waters Creates “Dead Zones”

Mid-Summer 2007



Read pages 556-558 in textbook

Area of mid-summer bottom-water hypoxia (< 2 mg O₂/L) In the Gulf of Mexico



Data source: Nancy N. Rabalais, UMMC, and R. Eugene Turner, LSU
Funding sources: NOAA Center for Sponsored Coastal Ocean Research and U.S. EPA Gulf of Mexico Program

Lectures 16-19

